

SmarcAL1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57715**Specification**

SmarcAL1 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC
Primary Accession	O9NZC9
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	105938

SmarcAL1 Polyclonal Antibody - Additional Information**Gene ID** 50485**Other Names**

SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A-like protein 1, 3.6.4.-, HepA-related protein, hHARP, Sucrose nonfermenting protein 2-like 1, SMARCAL1, HARP

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SmarcAL1 Polyclonal Antibody - Protein Information**Name** SMARCAL1**Synonyms** HARP**Function**

ATP-dependent annealing helicase that binds selectively to fork DNA relative to ssDNA or dsDNA and catalyzes the rewinding of the stably unwound DNA. Rewinds single-stranded DNA bubbles that are stably bound by replication protein A (RPA). Acts throughout the genome to reanneal stably unwound DNA, performing the opposite reaction of many enzymes, such as helicases and polymerases, that unwind DNA. May play an important role in DNA damage response by acting at stalled replication forks.

Cellular Location

Nucleus. Note=Recruited to damaged DNA regions

Tissue Location

Ubiquitously expressed, with high levels in testis.

SmarcAL1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SmarcAL1 Polyclonal Antibody - Images