

PPWD1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57740**Specification**

PPWD1 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q96BP3
Reactivity	Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	73575

PPWD1 Polyclonal Antibody - Additional Information**Gene ID** 23398**Other Names**

Peptidylprolyl isomerase domain and WD repeat-containing protein 1, 5.2.1.8, Spliceosome-associated cyclophilin, PPWD1 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=28954)
target="_blank">HGNC:28954)

Dilution

IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PPWD1 Polyclonal Antibody - Protein Information**Name** PPWD1 ([HGNC:28954](#))**Function**

PPlase that catalyzes the cis-trans isomerization of proline imidic peptide bonds in oligopeptides and may therefore assist protein folding (PubMed:20676357). May be involved in pre-mRNA splicing (PubMed:11991638).

Cellular Location

Nucleus. Note=Associated with spliceosomal complexes

PPWD1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PPWD1 Polyclonal Antibody - Images