

PABPC4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57747**Specification**

PABPC4 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q13310
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70783

PABPC4 Polyclonal Antibody - Additional Information**Gene ID** 8761**Other Names**

Polyadenylate-binding protein 4, PABP-4, Poly(A)-binding protein 4, Activated-platelet protein 1, APP-1, Inducible poly(A)-binding protein, iPABP, PABPC4, APP1, PABP4

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PABPC4 Polyclonal Antibody - Protein Information**Name** PABPC4**Synonyms** APP1, PABP4**Function**

Binds the poly(A) tail of mRNA (PubMed:8524242). Binds to SMIM26 mRNA and plays a role in its post-transcriptional regulation (PubMed:37009826). May be involved in cytoplasmic regulatory processes of mRNA metabolism. Can probably bind to cytoplasmic RNA sequences other than poly(A) in vivo (By similarity).

Cellular Location

Cytoplasm. Note=Localized in cytoplasmic mRNP granules containing untranslated mRNAs.

Tissue Location

Expressed at low levels in resting normal T cells; following T-cell activation, however, mRNA levels are rapidly up- regulated

PABPC4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PABPC4 Polyclonal Antibody - Images