

**Importin 8 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP57764****Specification**

---

**Importin 8 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	<a href="#">O15397</a>
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	119938

**Importin 8 Polyclonal Antibody - Additional Information****Gene ID** 10526**Other Names**

Importin-8, Imp8, Ran-binding protein 8, RanBP8, IPO8, RANBP8

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
ICC ~ ~ N/A  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Importin 8 Polyclonal Antibody - Protein Information****Name** IPO8**Synonyms** RANBP8**Function**

Involved in nuclear protein import, either by acting as autonomous nuclear transport receptor or as an adapter-like protein in association with the importin-beta subunit KPNB1. Acting autonomously, may serve as receptor for nuclear localization signals (NLS) and promote translocation of import substrates through the nuclear pore complex (NPC) by an energy requiring, Ran-dependent mechanism. At the nucleoplasmic side of the NPC, Ran binds to importin, the importin/substrate complex dissociates and importin is re-exported from the nucleus to the cytoplasm where GTP hydrolysis releases Ran. The directionality of nuclear import is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the

cytoplasm and nucleus (PubMed:<a href="http://www.uniprot.org/citations/9214382" target="\_blank">9214382</a>). In vitro mediates the nuclear import of the signal recognition particle protein SRP19 (PubMed:<a href="http://www.uniprot.org/citations/11682607" target="\_blank">11682607</a>). May also be involved in cytoplasm-to-nucleus shuttling of a broad spectrum of other cargos, including Argonaute- microRNAs complexes, the JUN protein, RELA/NF-kappa-B p65 subunit, the translation initiation factor EIF4E and a set of receptor-activated mothers against decapentaplegic homolog (SMAD) transcription factors that play a critical role downstream of the large family of transforming growth factor beta and bone morphogenetic protein (BMP) cytokines (Probable).

#### **Cellular Location**


Cytoplasm. Nucleus.

### **Importin 8 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Importin 8 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Importin 8) Polyclonal Antibody, Unconjugated (bs-20110R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.