

## **PIWIL1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57779

### **Specification**

# PIWIL1 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q96I94

Reactivity Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 99 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human PIWIL1

Epitope Specificity 421-520/861

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm. Note=Component of the

meiotic nuage, also named P granule, a germ-cell-specific organelle required to repress transposon during meiosis. Also

present in chromatoid body.

SIMILARITY Belongs to the argonaute family. Piwi

subfamily. Contains 1 PAZ domain.

Contains 1 Piwi domain.

SUBUNIT Interacts (via Piwi domain) with DICER1,

suggesting that it forms ribonucleoprotein

RISC complexes. This interaction is

regulated by HSP90AB1 activity. Interacts with MAEL, KIF17, PABPC1, PRMT5 and WDR77. Interacts (when methylated on

arginine residues) with TDRD1,

TDRKH/TDRD2, RNF17/TDRD4, TDRD6,

TDRD7 and TDRD9.

Post-translational modifications Arginine methylation by PRMT5 is required

for the interaction with Tudor domain-containing protein (TDRD1, TDRKH/TDRD2, RNF17/TDRD4, TDRD6, TDRD7 and TDRD9) and subsequent localization to the meiotic nuage, also

named P granule.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions** 

This gene encodes a member of the PIWI subfamily of Argonaute proteins, evolutionarily



conserved proteins containing both PAZ and Piwi motifs that play important roles in stem cell self-renewal, RNA silencing, and translational regulation in diverse organisms. The encoded protein may play a role as an intrinsic regulator of the self-renewal capacity of germline and hematopoietic stem cells. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2010].

## PIWIL1 Polyclonal Antibody - Additional Information

**Gene ID 9271** 

#### **Other Names**

Piwi-like protein 1, 3.1.26.-, PIWIL1

## Target/Specificity

Detected in most fetal and adult tissues. Expressed in testes, specifically in germline cells; detected in spermatocytes and spermatids during spermatogenesis. Increased expression in testicular tumors originating from embryonic germ cells with retention of germ cells phenotype. No expression in testicular tumors of somatic origin, such as Sertoli cell and Leydig cell tumors. Overexpressed in gastric cancer cells. Isoform 3 is ubiquitously expressed, and specifically in CD34+ hematopoietic progenitor cells but not in more differentiated cells.

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class ="dilution\_ICC">ICC~~N/A</span><br \> <span class ="dilution\_E">E~~N/A</span>

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

### **PIWIL1 Polyclonal Antibody - Protein Information**

### Name PIWIL1

#### **Function**

Endoribonuclease that plays a central role in postnatal germ cells by repressing transposable elements and preventing their mobilization, which is essential for the germline integrity. Acts via the piRNA metabolic process, which mediates the repression of transposable elements during meiosis by forming complexes composed of piRNAs and Piwi proteins and governs the methylation and subsequent repression of transposons. Directly binds methylated piRNAs, a class of 24 to 30 nucleotide RNAs that are generated by a Dicer-independent mechanism and are primarily derived from transposons and other repeated sequence elements. Strongly prefers a uridine in the first position of their guide (g1U preference, also named 1U-bias). Not involved in the piRNA amplification loop, also named ping-pong amplification cycle. Acts as an endoribonuclease that cleaves transposon messenger RNAs. Besides their function in transposable elements repression, piRNAs are probably involved in other processes during meiosis such as translation regulation. Probable component of some RISC complex, which mediates RNA cleavage and translational silencing. Also plays a role in the formation of chromatoid bodies and is required for some miRNAs stability. Required to sequester RNF8 in the cytoplasm until late spermatogenesis; RNF8 being released upon ubiquitination and degradation of PIWIL1.

# **Cellular Location**



Cytoplasm {ECO:0000250|UniProtKB:Q9JMB7}. Note=Component of the meiotic nuage, also named P granule, a germ-cell- specific organelle required to repress transposon activity during meiosis. Also present in chromatoid body {ECO:0000250|UniProtKB:Q9JMB7}

#### **Tissue Location**


Expressed in spermatocytes and spermatids. Also detected in prostate cancer (at protein level). Detected in most fetal and adult tissues. Expressed in testes, specifically in germline cells; detected in spermatocytes and spermatids during spermatogenesis Increased expression in testicular tumors originating from embryonic germ cells with retention of germ cells phenotype. No expression in testicular tumors of somatic origin, such as Sertoli cell and Leydig cell tumors. Overexpressed in gastric cancer cells. Isoform 3: Ubiquitously expressed, and specifically in CD34(+) hematopoietic progenitor cells but not in more differentiated cells

### **PIWIL1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **PIWIL1 Polyclonal Antibody - Images**




Blank control(blue): TM4 cells(fixed with 2% paraformaldehyde (10 min), then permeabilized with 90% ice-cold methanol for 30 min on ice).


Primary Antibody:Rabbit Anti-PIWIL1 antibody(bs-20224R), Dilution: 5  $\mu g$  in 100  $\mu L$  1X PBS containing 0.5% BSA;

Isotype Control Antibody: Rabbit IgG(orange) ,used under the same conditions );

Secondary Antibody: Goat anti-rabbit IgG-PE(white blue), Dilution: 1:200 in 1 X PBS containing 0.5% BSA.







Paraformaldehyde-fixed, paraffin embedded (Human stomach carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (PIWL1) Polyclonal Antibody, Unconjugated (bs-20224R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.