

Magphinin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57787**Specification**

Magphinin Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q12816
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	143716

Magphinin Polyclonal Antibody - Additional Information**Gene ID** 7216**Other Names**

Trophinin, MAGE-D3 antigen, TRO, KIAA1114, MAGED3

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Magphinin Polyclonal Antibody - Protein Information**Name** TRO**Synonyms** KIAA1114, MAGED3**Function**

Could be involved with bystin and tastin in a cell adhesion molecule complex that mediates an initial attachment of the blastocyst to uterine epithelial cells at the time of the embryo implantation. Directly responsible for homophilic cell adhesion.

Tissue Location

Strong expression at implantation sites. Found in the placenta from the sixth week of pregnancy. Was localized in the cytoplasm of the syncytiotrophoblast in the chorionic villi and in endometrial

decidual cells at the uteroplacental interface. After week 10, the level decreased and then disappeared from placental villi. Also found in macrophages

Magphinin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Magphinin Polyclonal Antibody - Images