

## **POP4 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP57839

### **Specification**

## **POP4 Polyclonal Antibody - Product Information**

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession <u>095707</u>

Reactivity
Host
Clonality
Calculated MW
Physical State

Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
25 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human POP4

Epitope Specificity 81-180/220

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Nucleus; nucleolus.
SIMILARITY Belongs to the eukaryotic

IMILARITY Belongs to the eukaryotic/archaeal RNase

P protein component 1 family.

SUBUNIT Component of nuclear RNase P and RNase

MRP ribonucleoproteins. RNase P consists of an RNA moiety and at least 8 protein subunits; POP1, RPP14, RPP20/POP7, RPP25, RPP29/POP4, RPP30, RPP38 and RPP40. RNase MRP consists of an RNA moiety and at least 9 protein subunits;

POP1, RPP14, RPP20/POP7, RPP25, RPP29/POP4, RPP30, RPP38, RPP40, POP5

and RPP21. Interacts with RPP25 and

POP5.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

# **Background Descriptions**

affinity purified by Protein A

This gene encodes one of the protein subunits of the small nucleolar ribonucleoprotein complexes: the endoribonuclease for mitochondrial RNA processing complex and the ribonuclease P complex. The encoded protein is localized to the nucleus and associates directly with the RNA component of these complexes. This protein is involved in processing of precursor RNAs. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Mar 2009]

#### POP4 Polyclonal Antibody - Additional Information

**Gene ID 10775** 



#### **Other Names**

Ribonuclease P protein subunit p29, hPOP4, 3.1.26.5, POP4, RPP29

#### **Dilution**

<span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \> <span class
="dilution\_IHC-F">IHC-F~~N/A</span><br \> <span class
="dilution\_IF">IF~~1:50~200</span><br \> <span class ="dilution\_ICC">ICC~~N/A</span><br \> <span class ="dilution\_E">E~~N/A</span>

#### **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

## **Storage**

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

# **POP4 Polyclonal Antibody - Protein Information**

#### Name POP4

Synonyms RPP29

#### **Function**

Component of ribonuclease P, a ribonucleoprotein complex that generates mature tRNA molecules by cleaving their 5'-ends.

#### **Cellular Location**

Nucleus, nucleolus.

### **POP4 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **POP4 Polyclonal Antibody - Images**