

RANBP16/Exp7 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57903**Specification**

RANBP16/Exp7 Polyclonal Antibody - Product Information

Application	IHC-P, WB
Primary Accession	O9UIA9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	123907

RANBP16/Exp7 Polyclonal Antibody - Additional Information**Gene ID** 23039**Other Names**

Exportin-7, Exp7, Ran-binding protein 16, XPO7, KIAA0745, RANBP16

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RANBP16/Exp7 Polyclonal Antibody - Protein Information**Name** XPO7**Synonyms** KIAA0745, RANBP16**Function**

Mediates the nuclear export of proteins (cargos) with broad substrate specificity. In the nucleus binds cooperatively to its cargo and to the GTPase Ran in its active GTP-bound form. Docking of this trimeric complex to the nuclear pore complex (NPC) is mediated through binding to nucleoporins. Upon transit of a nuclear export complex into the cytoplasm, disassembling of the complex and hydrolysis of Ran-GTP to Ran-GDP (induced by RANBP1 and RANGAP1, respectively) cause release of the cargo from the export receptor. XPO7 then return to the nuclear compartment and mediate another round of transport. The directionality of nuclear export is thought to be conferred by an asymmetric distribution of the GTP- and GDP-bound forms of Ran between the cytoplasm and nucleus.

Cellular Location

Cytoplasm. Nucleus Note=Shuttles between the nucleus and the cytoplasm

Tissue Location


Strong expression in testis, thyroid and bone marrow, low expression in lung, liver and small intestine, no expression in thymus, and remaining tissues studied have moderate expression. Expressed in red blood cells; overexpressed in red blood cells (cytoplasm) of patients with hereditary non-spherocytic hemolytic anemia of unknown etiology.

RANBP16/Exp7 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RANBP16/Exp7 Polyclonal Antibody - Images




Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RANBP16/Exp7) Polyclonal Antibody, Unconjugated (bs-21151R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20

minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RANBP16/Exp7) Polyclonal Antibody, Unconjugated (bs-21151R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:


A431(Human) Cell Lysate at 30 ug

Primary: Anti-Exp7 (bs-21151R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 124 kD

Observed band size: 124 kD



Sample:

Testis (Mouse) Lysate at 40 ug

Primary: Anti- RANBP16/Exp7 (bs-21151R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 124 kD

Observed band size: 124 kD