

SLC29A3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57946**Specification**

SLC29A3 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, ICC
Primary Accession	O9BZD2
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51815

SLC29A3 Polyclonal Antibody - Additional Information**Gene ID** 55315**Other Names**

Equilibrative nucleoside transporter 3, hENT3, Solute carrier family 29 member 3, SLC29A3, ENT3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SLC29A3 Polyclonal Antibody - Protein Information**Name** SLC29A3 ([HGNC:23096](#))**Synonyms** ENT3**Function**

Uniporter that mediates the facilitative transport of nucleoside across lysosomal and mitochondrial membranes (PubMed: [15701636](http://www.uniprot.org/citations/15701636), PubMed: [19164483](http://www.uniprot.org/citations/19164483), PubMed: [20595384](http://www.uniprot.org/citations/20595384), PubMed: [28729424](http://www.uniprot.org/citations/28729424)). Functions as a non-electrogenic Na(+)-independent transporter (PubMed: [15701636](http://www.uniprot.org/citations/15701636), PubMed: [19164483](http://www.uniprot.org/citations/19164483), PubMed: [28729424](http://www.uniprot.org/citations/28729424)). Substrate transport is pH-dependent and enhanced under acidic condition, probably reflecting the location of the transporter in acidic intracellular compartments (PubMed: [15701636](http://www.uniprot.org/citations/15701636), PubMed: [19164483](http://www.uniprot.org/citations/19164483), PubMed: [28729424](http://www.uniprot.org/citations/28729424)). Proton is not a cotransporting ion but most likely change the ionization state of the transporter which dictates

transport- permissible/impermissible conformation for nucleoside translocation (PubMed:28729424). May direct the nucleoside transport from lysosomes to cytosol or cytosol to mitochondria to facilitate the fundamental function of salvage synthesis of nucleic acids (PubMed:28729424). Involved in the transport of nucleosides (adenosine, guanosine, uridine, thymidine, cytidine and inosine) and deoxynucleosides (deoxyadenosine, deoxycytidine) (PubMed:15701636, PubMed:19164483, PubMed:20595384, PubMed:28729424). Also mediates transport of purine nucleobases (adenine, guanine) and pyrimidine nucleobases (uracil) (PubMed:15701636, PubMed:19164483). Also able to transport monoamine neurotransmitters dopamine, serotonin, noradrenaline and tyramine (PubMed:19164483). Capable of transporting ATP (PubMed:19164483). Mediates nucleoside export from lysosomes in macrophages, which regulates macrophage functions and numbers (By similarity).

Cellular Location

Lysosome membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Mitochondrion membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein. Note=Observed in a punctate intracellular pattern showing partial colocalization with late endosomes/lysosomes (PubMed:15701636). Detected at the cell surface only in certain placental cells (PubMed:19164483)

Tissue Location

Widely expressed in both adult and fetal tissues (PubMed:15701636). Highest levels in placenta, uterus, ovary, spleen, lymph node and bone marrow (PubMed:15701636). Expressed in liver (PubMed:19164483). Lowest levels in brain and heart (PubMed:15701636) Expressed in macrophages (PubMed:22174130)

SLC29A3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC29A3 Polyclonal Antibody - Images