

SLC35B4 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57951**Specification**

SLC35B4 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q969S0
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	37424

SLC35B4 Polyclonal Antibody - Additional Information**Gene ID** 84912**Other Names**

UDP-xylose and UDP-N-acetylglucosamine transporter, Solute carrier family 35 member B4, YEA4 homolog, SLC35B4, YEA4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SLC35B4 Polyclonal Antibody - Protein Information**Name** SLC35B4**Synonyms** YEA4**Function**

Antiporter that transports nucleotide sugars across the endoplasmic reticulum (ER) membrane in exchange for another nucleotide sugar. May couple UDP-alpha-D-glucuronate (UDP-GlcA) or UDP-alpha-D- xylose (UDP-Xyl) efflux to UDP-alpha-D-glucuronate (UDP-GlcA) influx into the ER lumen, which in turn stimulates glucuronidation and excretion of endobiotics and xenobiotics.

Cellular Location


Endoplasmic reticulum membrane; Multi-pass membrane protein

SLC35B4 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SLC35B4 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLC35B4) Polyclonal Antibody, Unconjugated (bs-21260R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SLC35B4) Polyclonal Antibody, Unconjugated (bs-21260R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.