

Spindly Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP57997

Specification

Spindly Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, ICC, E
Primary Accession	Q96EA4
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Protein Spindly
Epitope Specificity	401-500/605
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm, cytoskeleton, centrosome. Chromosome, centromere, kinetochore. Nucleus. Cytoplasm, cytoskeleton, spindle pole. Note=Localizes to the nucleus in interphase and to the kinetochore in early prometaphase. Relocalizes to the mitotic spindle pole before metaphase and is subsequently lost from the spindle poles after chromosome congression is completed. Removal of this protein from the kinetochore requires the dynein/dynactin complex. Belongs to the Spindly family. Phosphorylated upon DNA damage, probably by ATM or ATR.
SIMILARITY	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.
Post-translational modifications	
Important Note	

Background Descriptions

The five integral families of plant hormones consists of auxins, cytokinins, giberellins (GAs), abscisic acid (ABA), and ethylene. Giberellins, which consist of over fifty family members, mediate shoot growth. In *Arabidopsis*, SPINDLY (SPY) negatively regulates GA signal transduction. ERA1 (enhanced response to abscisic acid), which is identical to WIGGUM, controls floral and shoot apical meristem size and floral organ number in response to ABA. Ethylene is perceived by a family of five receptors, one of which is ETR1, whereas CTR1 is a negative regulator of the ethylene signal transduction pathway . Ethylene is also produced endogenously in *Arabidopsis* via a biosynthetic pathway, which is catalyzed by ACC synthase and ACC oxidase.

Spindly Polyclonal Antibody - Additional Information

Gene ID 54908

Other Names

Protein Spindly {ECO:0000255|HAMAP-Rule:MF_03041}, hSpindly, Arsenite-related gene 1 protein, Coiled-coil domain-containing protein 99 {ECO:0000255|HAMAP-Rule:MF_03041}, Rhabdomyosarcoma antigen MU-RMS-40.4A, Spindle apparatus coiled-coil domain-containing protein 1 {ECO:0000255|HAMAP-Rule:MF_03041}, SPDL1 {ECO:0000255|HAMAP-Rule:MF_03041}

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
ICC~~N/A
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Spindly Polyclonal Antibody - Protein Information

Name SPDL1 {ECO:0000255|HAMAP-Rule:MF_03041}

Function

Required for the localization of dynein and dynactin to the mitotic kintochore. Dynein is believed to control the initial lateral interaction between the kinetochore and spindle microtubules and to facilitate the subsequent formation of end-on kinetochore-microtubule attachments mediated by the NDC80 complex. Also required for correct spindle orientation. Does not appear to be required for the removal of spindle assembly checkpoint (SAC) proteins from the kinetochore upon bipolar spindle attachment (PubMed:17576797, PubMed:19468067). Acts as an adapter protein linking the dynein motor complex to various cargos and converts dynein from a non-processive to a highly processive motor in the presence of dynactin. Facilitates the interaction between dynein and dynactin and activates dynein processivity (the ability to move along a microtubule for a long distance without falling off the track) (PubMed:25035494). Plays a role in cell migration (PubMed:30258100).

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Chromosome, centromere, kinetochore. Nucleus Cytoplasm, cytoskeleton, spindle pole. Note=Localizes to the nucleus in interphase and to the kinetochore in early prometaphase. Relocalizes to the mitotic spindle pole before metaphase and is subsequently lost from the spindle poles after chromosome congression is completed. Removal of this protein from the kinetochore requires the dynein/dynactin complex

Spindly Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Spindly Polyclonal Antibody - Images

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Spindly Polyclonal Antibody, Unconjugated(bs-2321R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;
Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Spindly Polyclonal Antibody, Unconjugated(bs-2321R) 1:500, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining