

**FABP5 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58006****Specification****FABP5 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">Q01469</a>
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	15164

**FABP5 Polyclonal Antibody - Additional Information****Gene ID** 2171**Other Names**

Fatty acid-binding protein 5, Epidermal-type fatty acid-binding protein, E-FABP, Fatty acid-binding protein, epidermal, Psoriasis-associated fatty acid-binding protein homolog, PA-FABP, FABP5 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=3560](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=3560))  
HGNC:3560

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**FABP5 Polyclonal Antibody - Protein Information****Name** FABP5 ([HGNC:3560](#))**Function**

Intracellular carrier for long-chain fatty acids and related active lipids, such as endocannabinoids, that regulate the metabolism and actions of the ligands they bind. In addition to the cytosolic transport, selectively delivers specific fatty acids from the cytosol to the nucleus, wherein they activate nuclear receptors (PubMed: [22170058](http://www.uniprot.org/citations/22170058), PubMed: [21395585](http://www.uniprot.org/citations/21395585)). Delivers retinoic acid to the nuclear receptor peroxisome proliferator-activated receptor delta; which promotes proliferation and survival. May also serve as a synaptic carrier of endocannabinoid at central synapses and thus controls retrograde endocannabinoid signaling. Modulates inflammation by regulating PTGES induction via NF-kappa-B activation, and prostaglandin E2 (PGE2) biosynthesis during inflammation (By similarity). May be involved in keratinocyte differentiation (PubMed: [8092987](http://www.uniprot.org/citations/8092987)).

**Cellular Location**

Cytoplasm. Nucleus Synapse {ECO:0000250|UniProtKB:Q05816}. Postsynaptic density {ECO:0000250|UniProtKB:Q05816}. Secreted {ECO:0000250|UniProtKB:Q05816}. Note=Localizes primarily to the cytoplasm. Upon certain ligand binding, a conformation change exposes a nuclear localization motif and the protein is transported into nucleus (PubMed:24692551). Secreted by astrocytes, but not by neurons (By similarity). {ECO:0000250|UniProtKB:Q05816, ECO:0000269|PubMed:24692551}


**Tissue Location**

Keratinocytes; highly expressed in psoriatic skin (PubMed:8092987). Expressed in brain gray matter (PubMed:21395585)


**FABP5 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**FABP5 Polyclonal Antibody - Images**

Paraformaldehyde-fixed, paraffin embedded (mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FABP5) Polyclonal Antibody, Unconjugated (bs-2370R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (mouse stomach); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (FABP5) Polyclonal Antibody, Unconjugated (bs-2370R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.