

Urocortin 3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58057**Specification****Urocortin 3 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q969E3
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	4/18 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human UCN3
Epitope Specificity	120-157/161
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the sauvagine/corticotropin-releasing factor/urotensin I family.
SUBUNIT	Binds with high affinity to CRF receptors 2-alpha and 2-beta.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Urocortin 3 is a member of the sauvagine/corticotropin-releasing factor/urotensin I family. It is structurally related to the corticotropin-releasing factor (CRF) gene and the encoded product is an endogenous ligand for CRF type 2 receptors. In the brain it may be responsible for the effects of stress on appetite. In spite of the gene family name similarity, the product of this gene has no sequence similarity to urotensin II. A powerful cytoprotective effect of urocortin peptide administration against ischemia and reperfusion injury has been demonstrated in isolated cardiomyocyte models, as well as in the intact heart both in vitro and in vivo.

Urocortin 3 Polyclonal Antibody - Additional Information**Gene ID** 114131**Other Names**

Urocortin-3, Stresscopin, Urocortin III, Ucn III, UCN3, SPC

Dilution

IHC-P~~N/A<br \><span class

=["dilution_IHC-F">IHC-F](#)~[N/A](#)<br \><span class
=["dilution_IF">IF](#)~[1:50~200](#)<br \>E](#)~[N/A](#)

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Urocortin 3 Polyclonal Antibody - Protein Information

Name UCN3

Synonyms SPC

Function

Suppresses food intake, delays gastric emptying and decreases heat-induced edema. Might represent an endogenous ligand for maintaining homeostasis after stress.

Cellular Location


Secreted.

Urocortin 3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Urocortin 3 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Urocortin 3) Polyclonal Antibody, Unconjugated (bs-2786R) at 1:400 overnight at 4°C, followed by

operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.