

ABCE1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58064**Specification**

ABCE1 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P61221
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	67 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human ABCE1
Epitope Specificity	331-430/599
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Cytoplasm. Mitochondrion. Note=Localized to clusters of virus formation at the plasma membrane.
SIMILARITY	Belongs to the ABC transporter superfamily. ABCE family.Contains 2 4Fe-4S ferredoxin-type domains.Contains 2 ABC transporter domains.
SUBUNIT	Probably heterodimerizes with RNASEL; this interaction inhibits the RNASEL. Associates with HIV-1 Vif and HIV-1, HIV-2 and SIV Gag proteins.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

ABCE1 is a member of the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the OABP subfamily. It is alternatively referred to as the RNase L inhibitor as it functions to block the activity of ribonuclease L(referenced from entrez gene).

ABCE1 Polyclonal Antibody - Additional Information**Gene ID** 6059**Other Names**

ATP-binding cassette sub-family E member 1, 2'-5'-oligoadenylate-binding protein, HuHP68, RNase

L inhibitor, Ribonuclease 4 inhibitor, RNS4I, ABCE1, RLI, RNASEL1, RNASELI, RNS4I

Dilution

WB~1:1000
IHC-P~N/A
IHC-F~N/A
IF~1:50~200
E~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ABCE1 Polyclonal Antibody - Protein Information

Name ABCE1

Synonyms RLI, RNASEL1, RNASELI, RNS4I

Function

Nucleoside-triphosphatase (NTPase) involved in ribosome recycling by mediating ribosome disassembly (PubMed: [20122402](http://www.uniprot.org/citations/20122402)), PubMed: [21448132](http://www.uniprot.org/citations/21448132)). Able to hydrolyze ATP, GTP, UTP and CTP (PubMed: [20122402](http://www.uniprot.org/citations/20122402)). Splits ribosomes into free 60S subunits and tRNA- and mRNA-bound 40S subunits (PubMed: [20122402](http://www.uniprot.org/citations/20122402), PubMed: [21448132](http://www.uniprot.org/citations/21448132)). Acts either after canonical termination facilitated by release factors (ETF1/eRF1) or after recognition of stalled and vacant ribosomes by mRNA surveillance factors (PELO/Pelota) (PubMed: [20122402](http://www.uniprot.org/citations/20122402), PubMed: [21448132](http://www.uniprot.org/citations/21448132)). Involved in the No-Go Decay (NGD) pathway: recruited to stalled ribosomes by the Pelota-HBS1L complex, and drives the disassembly of stalled ribosomes, followed by degradation of damaged mRNAs as part of the NGD pathway (PubMed: [21448132](http://www.uniprot.org/citations/21448132)). Also plays a role in quality control of translation of mitochondrial outer membrane- localized mRNA (PubMed: [29861391](http://www.uniprot.org/citations/29861391)). As part of the PINK1-regulated signaling, ubiquitinated by CNOT4 upon mitochondria damage; this modification generates polyubiquitin signals that recruit autophagy receptors to the mitochondrial outer membrane and initiate mitophagy (PubMed: [29861391](http://www.uniprot.org/citations/29861391)). RNASEL-specific protein inhibitor which antagonizes the binding of 2-5A (5'-phosphorylated 2',5'-linked oligoadenylates) to RNASEL (PubMed: [9660177](http://www.uniprot.org/citations/9660177)). Negative regulator of the anti-viral effect of the interferon-regulated 2-5A/RNASEL pathway (PubMed: [11585831](http://www.uniprot.org/citations/11585831), PubMed: [9660177](http://www.uniprot.org/citations/9660177), PubMed: [9847332](http://www.uniprot.org/citations/9847332)).

Cellular Location


Cytoplasm. Mitochondrion

ABCE1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ABCE1 Polyclonal Antibody - Images



Sample:

Brain (Rat) Lysate at 40 ug

Liver (Rat) Lysate at 40 ug

Primary: Anti-ABCE1 (bs-2948R) at 1/300 dilution

Secondary: HRP conjugated Goat-Anti-rabbit IgG (bs-0295G-HRP) at 1/5000 dilution

Predicted band size: 67 kD

Observed band size: 66 kD