

CACNB2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58070

Specification

Physical State

CACNB2 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Rat, Dog
Host
Clonality
Polyclonal
Calculated MW
74 KDa

Immunogen KLH conjugated synthetic peptide derived

Liquid

from human CACNB2

Epitope Specificity 551-655/655

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane, sarcolemma; Peripheral

SUBUNIT

membrane protein; Cytoplasmic side.

Belongs to the calcium channel beta
subunit family.Contains 1 SH3 domain.

The L-type calcium channel is composed.

The L-type calcium channel is composed of four subunits: alpha-1, alpha-2, beta and gamma. Interacts with RRAD. Interaction with RRAD regulates the trafficking of

CACNA1C to the cell membrane.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

This gene encodes a subunit of a voltage-dependent calcium channel protein that is a member of the voltage-gated calcium channel superfamily. The gene product was originally identified as an antigen target in Lambert-Eaton myasthenic syndrome, an autoimmune disorder. Mutations in this gene are associated with Brugada syndrome. Alternatively spliced variants encoding different isoforms have been described. [provided by RefSeq, Feb 2013]

CACNB2 Polyclonal Antibody - Additional Information

Gene ID 783

Other Names

Voltage-dependent L-type calcium channel subunit beta-2, CAB2, Calcium channel voltage-dependent subunit beta 2, Lambert-Eaton myasthenic syndrome antigen B, MYSB, CACNB2, CACNLB2, MYSB



Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

CACNB2 Polyclonal Antibody - Protein Information

Name CACNB2

Synonyms CACNLB2, MYSB

Function

Beta subunit of voltage-dependent calcium channels which contributes to the function of the calcium channel by increasing peak calcium current (By similarity). Plays a role in shifting voltage dependencies of activation and inactivation of the channel (By similarity). May modulate G protein inhibition (By similarity). May contribute to beta-adrenergic augmentation of Ca(2+) influx in cardiomyocytes, thereby regulating increases in heart rate and contractile force (PubMed:36424916). Involved in membrane targeting of the alpha-1 subunit CACNA1C (PubMed:17525370).

Cellular Location

Cell membrane, sarcolemma; Peripheral membrane protein; Cytoplasmic side

Tissue Location

Expressed in all tissues.

CACNB2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

CACNB2 Polyclonal Antibody - Images