

**MTF-1 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58078**

**Specification**

**MTF-1 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q14872</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	81 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human MTF-1
Epitope Specificity	101-200/753
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Contains 6 C2H2-type zinc fingers.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

The zinc finger transcription factor MTF-1 (metal-responsive transcription factor-1) is conserved from insects to vertebrates. The major role of MTF-1 in both organisms is to control the transcription of genes involved in the homeostasis and detoxification of heavy metal ions such as Cu<sup>2+</sup>, Zn<sup>2+</sup> and Cd<sup>2+</sup>. In mammals, MTF-1 serves at least two additional roles. First, targeted disruption of the MTF-1 gene results in death at embryonic day 14 due to liver degeneration, revealing a stage-specific developmental role. Second, under hypoxic-anoxic stress, MTF-1 helps to activate the transcription of the gene placental growth factor (PIGF), an angiogenic protein.

**MTF-1 Polyclonal Antibody - Additional Information**

**Gene ID** 4520

**Other Names**

Metal regulatory transcription factor 1, MRE-binding transcription factor, Transcription factor MTF-1, MTF1

**Dilution**

IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
E~N/A

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**MTF-1 Polyclonal Antibody - Protein Information**

**Name** MTF1

**Function**

Zinc-dependent transcriptional regulator of cellular adaption to conditions of exposure to heavy metals (PubMed:<a href="http://www.uniprot.org/citations/8065932" target="\_blank">8065932</a>). Binds to metal responsive elements (MRE) in promoters and activates the transcription of metallothionein genes like metallothionein-2/MT2A (PubMed:<a href="http://www.uniprot.org/citations/8065932" target="\_blank">8065932</a>). Also regulates the expression of metalloproteases in response to intracellular zinc and functions as a catabolic regulator of cartilages (By similarity).


**Cellular Location**

Nucleus. Cytoplasm {ECO:0000250|UniProtKB:Q07243}. Note=Translocation to the nucleus is induced by metals. {ECO:0000250|UniProtKB:Q07243}

**MTF-1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MTF-1 Polyclonal Antibody - Images**

Tissue/cell: human gastric carcinoma; 4% Paraformaldehyde-fixed and paraffin-embedded;  
Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at

37°C for 20 min;

Incubation: Anti-MTF-1 Polyclonal Antibody, Unconjugated(bs-3601R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining