

DACH2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58087

Specification

DACH2 Polyclonal Antibody - Product Information

| | |
|--------------------------------|---|
| Application | WB, IHC-P, IHC-F, IF, E |
| Primary Accession | O96NX9 |
| Reactivity | Rat, Pig, Dog, Bovine |
| Host | Rabbit |
| Clonality | Polyclonal |
| Calculated MW | 65 KDa |
| Physical State | Liquid |
| Immunogen | KLH conjugated synthetic peptide derived from human DACH2 |
| Epitope Specificity | 501-599/599 |
| Isotype | IgG |
| Purity | |
| affinity purified by Protein A | |
| Buffer | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol. |
| SUBCELLULAR LOCATION | Nuclear. |
| SIMILARITY | Belongs to the DACH/dachshund family. |
| SUBUNIT | Interacts with SIX6 and EYA2. |
| Important Note | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

Background Descriptions

DACH2 is a transcription factor that is involved in regulation of organogenesis. It seems to be a regulator for SIX1 and SIX6 and may act as a corepressor of SIX6 in regulating proliferation by directly repressing cyclin-dependent kinase inhibitors, including the p27Kip1 promoter. It probably binds to DNA via its DACH box-N domain.

DACH2 Polyclonal Antibody - Additional Information

Gene ID 117154

Other Names

Dachshund homolog 2, Dach2, DACH2

Dilution

WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>E~~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

DACH2 Polyclonal Antibody - Protein Information**Name** DACH2**Function**

Transcription factor that is involved in regulation of organogenesis. Seems to be a regulator for SIX1 and SIX6. Seems to act as a corepressor of SIX6 in regulating proliferation by directly repressing cyclin-dependent kinase inhibitors, including the p27Kip1 promoter. Is recruited with SIX6 to the p27Kip1 promoter in embryonal retina. SIX6 corepression also seems to involve NCOR1, TBL1, HDAC1 and HDAC3. May be involved together with PAX3, SIX1, and EYA2 in regulation of myogenesis. In the developing somite, expression of DACH2 and PAX3 is regulated by the overlying ectoderm, and DACH2 and PAX3 positively regulate each other's expression (By similarity). Probably binds to DNA via its DACHbox-N domain.

Cellular Location

Nucleus.

DACH2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DACH2 Polyclonal Antibody - Images