

**Neurogenin 2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58092****Specification**

---

**Neurogenin 2 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q9H2A3</a>
Reactivity	Rat, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28621

**Neurogenin 2 Polyclonal Antibody - Additional Information****Gene ID** 63973**Other Names**

Neurogenin-2, NGN-2, Class A basic helix-loop-helix protein 8, bHLHa8, Protein atonal homolog 4, NEUROG2, ATOH4, BHLHA8, NGN2

**Dilution**

WB~1:1000  
IHC-P~N/A  
IHC-F~N/A  
IF~1:50~200  
E~N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Neurogenin 2 Polyclonal Antibody - Protein Information****Name** NEUROG2**Synonyms** ATOH4, BHLHA8, NGN2**Function**

Transcriptional regulator. Involved in neuronal differentiation. Activates transcription by binding to the E box (5'- CANNTG-3').

**Cellular Location**


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981}.

## Neurogenin 2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Neurogenin 2 Polyclonal Antibody - Images




Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Neurogenin 2) Polyclonal Antibody, Unconjugated (bs-3692R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-Neurogenin 2 Polyclonal Antibody, Unconjugated(bs-3692R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



**Sample:**

Cerebrum (Mouse) Lysate at 40 ug

Primary: Anti-Neurogenin 2 (bs-3692R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 30 kD

Observed band size: 30 kD