

# **Protor-1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58103

## **Specification**

# **Protor-1 Polyclonal Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IHC-P, IHC-F, IF
P85299
Rat, Pig, Dog, Bovine
Rabbit
Polyclonal
42753

# **Protor-1 Polyclonal Antibody - Additional Information**

**Gene ID** 55615

#### **Other Names**

Proline-rich protein 5, Protein observed with Rictor-1, Protor-1, PRR5, PROTOR1

## **Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

#### Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **Protor-1 Polyclonal Antibody - Protein Information**

Name PRR5

# Synonyms PROTOR1

# **Function**

Subunit of mTORC2, which regulates cell growth and survival in response to hormonal signals. mTORC2 is activated by growth factors, but, in contrast to mTORC1, seems to be nutrient-insensitive. mTORC2 seems to function upstream of Rho GTPases to regulate the actin cytoskeleton, probably by activating one or more Rho-type guanine nucleotide exchange factors. mTORC2 promotes the serum-induced formation of stress-fibers or F-actin. mTORC2 plays a critical role in AKT1 'Ser-473' phosphorylation, which may facilitate the phosphorylation of the activation loop of AKT1 on 'Thr-308' by PDK1 which is a prerequisite for full activation. mTORC2 regulates the phosphorylation of SGK1 at 'Ser-422'. mTORC2 also modulates the phosphorylation of PRKCA on 'Ser-657'. PRR5 plays an important role in regulation of PDGFRB expression and in modulation of platelet-derived growth factor signaling. May act as a tumor suppressor in breast cancer.

### **Tissue Location**

Most abundant in kidney and liver. Also highly expressed in brain, spleen, testis and placenta. Overexpressed in several colorectal tumors.





101. 000.070.1000 1 ax. 000.070.100

# **Protor-1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Protor-1 Polyclonal Antibody - Images