

ZMIZ2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58183**Specification**

ZMIZ2 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q8NF64
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	96537

ZMIZ2 Polyclonal Antibody - Additional Information**Gene ID** 83637**Other Names**

Zinc finger MIZ domain-containing protein 2, PIAS-like protein Zimp7, ZMIZ2, KIAA1886, ZIMP7

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ZMIZ2 Polyclonal Antibody - Protein Information**Name** ZMIZ2**Synonyms** KIAA1886, ZIMP7**Function**

Increases ligand-dependent transcriptional activity of AR and other nuclear hormone receptors.

Cellular Location

Nucleus. Note=Detected at replication foci throughout S phase

Tissue Location


Expressed most abundantly in testis with lower levels in heart, brain, pancreas, prostate and ovary

ZMIZ2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZMIZ2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZMIZ2) Polyclonal Antibody, Unconjugated (bs-4391R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.