

CUEDC2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58237**Specification**

CUEDC2 Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q9H467
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	32009

CUEDC2 Polyclonal Antibody - Additional Information**Gene ID** 79004**Other Names**

CUE domain-containing protein 2, CUEDC2, C10orf66

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CUEDC2 Polyclonal Antibody - Protein Information**Name** CUEDC2**Synonyms** C10orf66**Function**

Down-regulates ESR1 protein levels through the ubiquitination-proteasome pathway, regardless of the presence of 17 beta-estradiol. Also involved in 17 beta-estradiol-induced ESR1 degradation. Controls PGR protein levels through a similar mechanism.

Cellular Location

Cytoplasm. Nucleus

Tissue Location


Significantly up-regulated in breast tumor tissues compared with matched adjacent normal tissues (at protein level) Levels inversely correlate with ESR1 in breast cancers and are lower in low-grade tumors compared to high-grade tumors

CUEDC2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CUEDC2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse lung tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CUEDC2) Polyclonal Antibody, Unconjugated (bs-4608R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.