

Apolipoprotein A V Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58266**Specification**

Apolipoprotein A V Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q6Q788
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	41213

Apolipoprotein A V Polyclonal Antibody - Additional Information**Gene ID** 116519**Other Names**

Apolipoprotein A-V, Apo-AV, ApoA-V, Apolipoprotein A5, Regeneration-associated protein 3, APOA5, RAP3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Apolipoprotein A V Polyclonal Antibody - Protein Information**Name** APOA5**Synonyms** RAP3**Function**

Minor apolipoprotein mainly associated with HDL and to a lesser extent with VLDL. May also be associated with chylomicrons. Important determinant of plasma triglyceride (TG) levels by both being a potent stimulator of apo-CII lipoprotein lipase (LPL) TG hydrolysis and an inhibitor of the hepatic VLDL-TG production rate (without affecting the VLDL-apoB production rate) (By similarity). Activates poorly lecithin:cholesterol acyltransferase (LCAT) and does not enhance efflux of cholesterol from macrophages. Binds heparin (PubMed:17326667).

Cellular Location

Secreted. Early endosome. Late endosome. Golgi apparatus, trans-Golgi network. Note=In the presence of SORL1, internalized to early endosomes, sorted in a retrograde fashion to late endosomes, from which a portion is sent to lysosomes and degradation, another portion is sorted to the trans-Golgi network


Tissue Location

Liver and plasma.

Apolipoprotein A V Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Apolipoprotein A V Polyclonal Antibody - Images

Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Apolipoprotein A V) Polyclonal Antibody, Unconjugated (bs-5035R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.