

**GDF10 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58299****Specification****GDF10 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">P55107</a>
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	12/53 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human GDF10
Epitope Specificity	371-478/478
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted.
SIMILARITY	Belongs to the TGF-beta family.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

GDF10, or growth differentiation factor 10, bone morphogenetic protein (BMP) family and the TGF-beta superfamily. This group of proteins is characterized by a polybasic proteolytic processing site which is cleaved to produce a mature protein containing seven conserved cysteine residues. GDF10 is expressed in femur, brain, lung, skeletal muscle, pancreas and testis.

**GDF10 Polyclonal Antibody - Additional Information****Gene ID** 2662**Other Names**

Growth/differentiation factor 10, GDF-10, Bone morphogenetic protein 3B, BMP-3B, Bone-inducing protein, BIP, GDF10 ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=4215](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=4215) target="\_blank">HGNC:4215</a>), BMP3B

**Target/Specificity**

Expressed in femur, brain, lung, skeletal muscle, pancreas and testis.

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class

=>IHC-F~N/A</span><br \><span class  
=>IF~1:50~200</span><br \><span class =>E~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**GDF10 Polyclonal Antibody - Protein Information**

**Name** GDF10 ([HGNC:4215](#))

**Synonyms** BMP3B

**Function**

Growth factor involved in osteogenesis and adipogenesis. Plays an inhibitory role in the process of osteoblast differentiation via SMAD2/3 pathway. Plays an inhibitory role in the process of adipogenesis.

**Cellular Location**

Secreted {ECO:0000250|UniProtKB:P97737}.


**Tissue Location**


Expressed in femur, brain, lung, skeletal muscle, pancreas and testis.

**GDF10 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**GDF10 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (GDF10) Polyclonal Antibody, Unconjugated (bs-5720R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.