

### **ADAMTS8 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58333

# **Specification**

### **ADAMTS8 Polyclonal Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity Isotype **Purity** affinity purified by Protein A

Buffer

SUBCELLULAR LOCATION

**SIMILARITY** 

Post-translational modifications

Important Note

**Background Descriptions** 

WB, IHC-P, IHC-F, IF, E

O9UP79
Rat, Pig, Dog
Rabbit
Polyclonal
74 KDa
Liquid

KLH conjugated synthetic peptide derived

from human ADAMTS8

541-640/889

laG

0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

Secreted, extracellular space, extracellular

matrix (By similarity).

Contains 1 disintegrin domain. Contains 1

peptidase M12B domain. Contains 2 TSP

type-1 domains.

The precursor is cleaved by a furing and appreciate (Py similarity)

endopeptidase (By similarity).

Glycosylated. Can be O-fucosylated by POFUT2 on a serine or a threonine residue found within the consensus sequence C1-X(2)-(S/T)-C2-G of the TSP type-1 repeat domains where C1 and C2 are the first and second cysteine residue of the repeat, respectively. Fucosylated repeats can then be further glycosylated by the addition of a beta-1,3-glucose residue by the

glucosyltransferase, B3GALTL.
Fucosylation mediates the efficient
secretion of ADAMTS family members. Also
can be C-glycosylated with one or two
mannose molecules on tryptophan
residues within the consensus sequence
W-X-X-W of the TPRs, and N-glycosylated.
These other glycosylations can also

facilitate secretion (By similarity). This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.



ADAMTS proteases are secreted enzymes containing a prometalloprotease domain of the reprolysin type. The ADAMTS proteases function in processing of procollagens and von Willebrand factor as well as catabolism of aggrecan, versican and brevican. They have been demonstrated to have important roles in connective tissue organization, coagulation, inflammation, arthritis, angiogenesis and cell migration. A member of the metalloproteinase family containing disintegrin like domains (ADAMs), the function of ADAMTS8 is still poorly understood. ADAMTS8 contains the canonical HExxHxxxxxH zinc metalloproteinase motif, and has been shown to be proteolytically active on a range of substrates. ADAMTS8 is inhibited by the endogenous MMP inhibitors, TIMP1, 2, 3 and 4, but most efficiently by TIMP3. In addition to the metalloprotease domain, ADAMTS8 has a propeptide domain, a Prohormone Convertase (PC, furin) cleavage site, a cysteine rich domain and thrombospondin 1 like domains.

# **ADAMTS8 Polyclonal Antibody - Additional Information**

**Gene ID** 11095

### **Other Names**

A disintegrin and metalloproteinase with thrombospondin motifs 8, ADAM-TS 8, ADAM-TS 8, ADAMTS-8, 3.4.24.-, METH-2, METH-8, ADAMTS8, METH2

## **Target/Specificity**

Highly expressed in adult and fetal lung, lower expression in brain, placenta, heart, stomach and fetal brain and kidney.

#### **Dilution**

```
<span class ="dilution_WB">WB~~1:1000</span><br \><span class
="dilution_IHC-P">IHC-P~~N/A</span><br \><span class
="dilution_IHC-F">IHC-F~~N/A</span><br \><span class
="dilution_IF">IF~~1:50~200</span><br \><span class ="dilution_E">E~~N/A</span>
```

#### Storage

Store at -20  $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4  $^{\circ}$ C.

### **ADAMTS8 Polyclonal Antibody - Protein Information**

Name ADAMTS8

**Synonyms** METH2

### **Function**

Has anti-angiogenic properties.

## **Cellular Location**

Secreted, extracellular space, extracellular matrix

## **Tissue Location**

Highly expressed in adult and fetal lung, lower expression in brain, placenta, heart, stomach and fetal brain and kidney

## **ADAMTS8 Polyclonal Antibody - Protocols**







Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

**ADAMTS8 Polyclonal Antibody - Images**