

**Hyaluronidase3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58341****Specification****Hyaluronidase3 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">O43820</a>
Reactivity	Rat, Pig
Host	Rabbit
Clonality	Polyclonal
Calculated MW	47 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human Hyaluronidase3
Epitope Specificity	51-150/417
Isotype	IgG
<b>Purity</b>	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Secreted (By similarity). Lysosome (By similarity).
SIMILARITY	Belongs to the glycosyl hydrolase 56 family. Contains 1 EGF-like domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

**Background Descriptions**

HYAL3 is a protein which is similar in structure to hyaluronidases. Hyaluronidases intracellularly degrade hyaluronan, one of the major glycosaminoglycans of the extracellular matrix. Hyaluronan is thought to be involved in cell proliferation, migration and differentiation. However, this protein has not yet been shown to have hyaluronidase activity.

**Hyaluronidase3 Polyclonal Antibody - Additional Information****Gene ID** 8372**Other Names**

Hyaluronidase-3, Hyal-3, 3.2.1.35, Hyaluronoglucosaminidase-3, Lung carcinoma protein 3, LuCa-3, HYAL3, LUCA3

**Target/Specificity**

Bone marrow, testis and kidney. Isoform 4 is detected in all bladder tumor and prostate tumor cells.

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class

="dilution\_IHC-P">IHC-P~N/A</span><br \><span class="dilution\_IHC-F">IHC-F~N/A</span><br \><span class="dilution\_IF">IF~1:50~200</span><br \><span class="dilution\_E">E~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**Hyaluronidase3 Polyclonal Antibody - Protein Information**

**Name** HYAL3

**Synonyms** LUCA3

**Function**

Facilitates sperm penetration into the layer of cumulus cells surrounding the egg by digesting hyaluronic acid. Involved in induction of the acrosome reaction in the sperm. Involved in follicular atresia, the breakdown of immature ovarian follicles that are not selected to ovulate. Induces ovarian granulosa cell apoptosis, possibly via apoptotic signaling pathway involving CASP8 and CASP3 activation, and poly(ADP-ribose) polymerase (PARP) cleavage. Has no hyaluronidase activity in embryonic fibroblasts in vitro. Has no hyaluronidase activity in granulosa cells in vitro.

**Cellular Location**

Secreted {ECO:0000250|UniProtKB:Q8VEI3}. Cell membrane {ECO:0000250|UniProtKB:Q8VEI3}. Cytoplasmic vesicle, secretory vesicle, acrosome {ECO:0000250|UniProtKB:Q8VEI3}. Endoplasmic reticulum {ECO:0000250|UniProtKB:Q8VEI3}. Early endosome {ECO:0000250|UniProtKB:Q8VEI3}. Note=Mostly present in low-density vesicles. Low levels in higher density vesicles of late endosomes and lysosomes. Localized in punctate cytoplasmic vesicles and in perinuclear structures, but does not colocalize with LAMP1. Localized on the plasma membrane over the acrosome and on the surface of the midpiece of the sperm tail. {ECO:0000250|UniProtKB:Q8VEI3}

**Tissue Location**


Expressed in sperm (PubMed:20586096). Highly expressed in epidermis of the skin, where it is expressed intracellularly in the deep horny layer (at protein level) (PubMed:21699545). Bone marrow, testis and kidney (PubMed:10493834)

**Hyaluronidase3 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**Hyaluronidase3 Polyclonal Antibody - Images**



**Sample:**


Mouse(Kidney) Lysate at 40 ug

Primary: Anti- Hyaluronidase3 (bs-5889R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 47 kD

Observed band size: 48 kD



Blank control:Molt4.

Primary Antibody (green line): Rabbit Anti-Hyaluronidase3 antibody (bs-5889R)

Dilution: 2 µg /10<sup>6</sup> cells;

Isotype Control Antibody (orange line): Rabbit IgG .

Secondary Antibody : Goat anti-rabbit IgG-PE

Dilution: 1 µg /test.

**Protocol**

The cells were incubated in 5%BSA to block non-specific protein-protein interactions for 30 min at room temperature .Cells stained with Primary Antibody for 30 min at room temperature. The secondary antibody used for 40 min at room temperature. Acquisition of 20,000 events was performed.