

SNF2L Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58362**Specification**

SNF2L Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	P28370
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	122605

SNF2L Polyclonal Antibody - Additional Information**Gene ID** 6594**Other Names**

Probable global transcription activator SNF2L1, 3.6.4.-, ATP-dependent helicase SMARCA1, Nucleosome-remodeling factor subunit SNF2L, SWI/SNF-related matrix-associated actin-dependent regulator of chromatin subfamily A member 1, SMARCA1, SNF2L, SNF2L1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SNF2L Polyclonal Antibody - Protein Information**Name** SMARCA1**Synonyms** SNF2L, SNF2L1**Function**

[Isoform 1]: Catalytically inactive when either DNA or nucleosomes are the substrate and does not possess chromatin-remodeling activity (PubMed:15310751, PubMed:28801535). Acts as a negative regulator of chromatin remodelers by generating inactive complexes (PubMed:15310751).

Cellular Location

Nucleus.

Tissue Location

[Isoform 1]: Mainly expressed in non-neuronal tissues such as lung, breast, kidney, and ovary

SNF2L Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SNF2L Polyclonal Antibody - Images