

METTL11A Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58368**Specification**

METTL11A Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q9BV86
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25387

METTL11A Polyclonal Antibody - Additional Information**Gene ID** 28989**Other Names**

N-terminal Xaa-Pro-Lys N-methyltransferase 1, 2.1.1.244, Alpha N-terminal protein methyltransferase 1A, Methyltransferase-like protein 11A, N-terminal RCC1 methyltransferase, X-Pro-Lys N-terminal protein methyltransferase 1A, NTM1A, N-terminal Xaa-Pro-Lys N-methyltransferase 1, N-terminally processed, NTMT1, C9orf32, METTL11A, NRMT, NRMT1

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

METTL11A Polyclonal Antibody - Protein Information**Name** NTMT1**Synonyms** C9orf32, METTL11A, NRMT, NRMT1**Function**

Distributive alpha-N-methyltransferase that methylates the N- terminus of target proteins containing the N-terminal motif [Ala/Gly/Pro/Ser]-Pro-Lys when the initiator Met is cleaved. Specifically catalyzes mono-, di- or tri-methylation of the exposed alpha-amino group of the Ala, Gly or Ser residue in the [Ala/Gly/Ser]- Pro-Lys motif and mono- or di-methylation of Pro in the Pro-Pro-Lys motif. Some of the substrates may be primed by NTMT2-mediated monomethylation (PubMed:24090352). Catalyzes the trimethylation of the N-terminal Gly in CENPA (after removal of Met-1). Responsible for the N-terminal methylation of KLHL31, MYL2, MYL3, RB1, RCC1, RPL23A and SET. Required during mitosis for normal bipolar spindle formation and chromosome segregation via its action on RCC1.


Cellular Location

Nucleus. Note=Predominantly nuclear (PubMed:24090352)

METTL11A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

METTL11A Polyclonal Antibody - Images

Tissue/cell: mouse embryo tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;


Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-METTL11A Polyclonal Antibody, Unconjugated(bs-6057R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining




Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20

minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (METTL11A) Polyclonal Antibody, Unconjugated (bs-6057R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (METTL11A) Polyclonal Antibody, Unconjugated (bs-6057R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 30min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (METTL11A) Polyclonal Antibody, Unconjugated (bs-6057R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.