

AKR1B10 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) **Catalog # AP58407**

Specification

AKR1B10 Polyclonal Antibody - Product Information

WB, IHC-P, IHC-F, IF, E Application

Primary Accession 060218 Reactivity Rat Host Rabbit Clonality **Polyclonal** Calculated MW **35 KDa Physical State**

Immunogen KLH conjugated synthetic peptide derived

Liquid

laG

from human AKR1B10

8-110/316 **Epitope Specificity**

Isotype **Purity** affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Lysosome. Secreted. Note=Secreted

through a lysosome-mediated non-classical

pathway.

SIMILARITY Belongs to the aldo/keto reductase family. Important Note This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

AKR1B10 is also known as aldose reductase-like-1 (ARL-1), small intestine reductase (SI reductase) or aldose reductase-related protein (ARP or hARP). AKR1B10 is found in many tissues but is predominantly expressed in small intestine, colon and adrenal gland. AKR1B10 is an efficient reductase for aliphatic and aromatic aldehydes. It plays a role in steroid metabolism as well as detoxification of aldehydes in digested food, and may be involved in the retinal-retinoic acid signaling pathway. AKR1B10 is prominently overexpressed in non-small cell lung carcinoma and adenocarcinoma. Cigarette smoking is an independent variable responsible for this overexpression. AKR1B10 may play a role regulating cell proliferation and cellular response to carbonyl stress.

AKR1B10 Polyclonal Antibody - Additional Information

Gene ID 57016

Other Names

Aldo-keto reductase family 1 member B10, 1.1.1.300, 1.1.1.54, ARL-1, Aldose reductase-like, Aldose reductase-related protein, ARP, hARP, Small intestine reductase, SI reductase, AKR1B10, AKR1B11



Target/Specificity

Found in many tissues. Highly expressed in small intestine, colon and adrenal gland.

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

AKR1B10 Polyclonal Antibody - Protein Information

Name AKR1B10

Synonyms AKR1B11

Function

Catalyzes the NADPH-dependent reduction of a wide variety of carbonyl-containing compounds to their corresponding alcohols (PubMed: 12732097, PubMed:18087047, PubMed:19013440, PubMed:19563777, PubMed:9565553). Displays strong enzymatic activity toward all-trans- retinal, 9-cis-retinal, and 13-cis-retinal (PubMed: 12732097, PubMed:18087047). Plays a critical role in detoxifying dietary and lipid-derived unsaturated carbonyls, such as crotonaldehyde, 4- hydroxynonenal, trans-2-hexenal, trans-2,4-hexadienal and their glutathione-conjugates carbonyls (GS-carbonyls) (PubMed:19013440, PubMed:19563777). Displays no reductase activity towards glucose (PubMed: 12732097).

Cellular Location

Lysosome. Secreted. Note=Secreted through a lysosome- mediated non-classical pathway

Tissue Location

Found in many tissues. Highly expressed in small intestine, colon and adrenal gland.

AKR1B10 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation



- Flow Cytomety
- Cell Culture

AKR1B10 Polyclonal Antibody - Images



Sample:

A549 Cell Lysate at 40 ug SW480 Cell Lysate at 40 ug

_x005f Primary: Anti- AKR1B10 (bs-6274R) at 1/300 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 35 kD Observed band size: 35 kD