

TKTL1 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58409**Specification**

TKTL1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P51854
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	65333

TKTL1 Polyclonal Antibody - Additional Information**Gene ID** 8277**Other Names**

Transketolase-like protein 1, 2.2.1.1, Transketolase 2, TK 2, Transketolase-related protein, TKTL1, TKR, TKT2

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TKTL1 Polyclonal Antibody - Protein Information**Name** TKTL1**Synonyms** TKR, TKT2**Function**

Catalyzes the transfer of a two-carbon ketol group from a ketose donor to an aldose acceptor, via a covalent intermediate with the cofactor thiamine pyrophosphate.

Cellular Location

Cytoplasm.

Tissue Location


Widely expressed (PubMed:8838793). Expressed in endothelial cells and in peripheral neurons (at protein level) (PubMed:15991799). [Isoform 4]: Expressed in fetal neocortex.

TKTL1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TKTL1 Polyclonal Antibody - Images



Sample:

293T(Human) Cell Lysate at 30 ug


A431(Human) Cell Lysate at 30 ug

Primary: Anti-TKTL1 (bs-6284R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 65 kD

Observed band size: 65 kD



Sample:


Testis (Mouse) Lysate at 40 ug

Primary: Anti-TKTL1 (bs-6284R) at 1/1000 dilution


Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 65 kD

Observed band size: 65 kD



Paraformaldehyde-fixed, paraffin embedded (Rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TKTL1) Polyclonal Antibody, Unconjugated (bs-6284R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse testis); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TKTL1) Polyclonal Antibody, Unconjugated (bs-6284R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.