

**MDFIC Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58431****Specification****MDFIC Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q9P1T7</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25788

**MDFIC Polyclonal Antibody - Additional Information****Gene ID** 29969**Other Names**

MyoD family inhibitor domain-containing protein, I-mfa domain-containing protein, hIC, MDFIC ([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=28870](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=28870))  
target="\_blank">HGNC:28870</a>)

**Dilution**

<span class = "dilution\_WB">WB~~1:1000</span><br \><span class = "dilution\_IHC-P">IHC-P~~N/A</span><br \><span class = "dilution\_IHC-F">IHC-F~~N/A</span><br \><span class = "dilution\_IF">IF~~1:50~200</span><br \><span class = "dilution\_E">E~~N/A</span>

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**MDFIC Polyclonal Antibody - Protein Information****Name** MDFIC ([HGNC:28870](#))**Function**

Required to control the activity of various transcription factors through their sequestration in the cytoplasm. Retains nuclear Zic proteins ZIC1, ZIC2 and ZIC3 in the cytoplasm and inhibits their transcriptional activation (By similarity). Modulates the expression from cellular promoters. Binds to the axin complex, resulting in an increase in the level of free beta-catenin (PubMed:<a href="http://www.uniprot.org/citations/12192039" target="\_blank">12192039</a>). Affects axin regulation of the WNT and JNK signaling pathways (PubMed:<a href="http://www.uniprot.org/citations/12192039" target="\_blank">12192039</a>). Involved in the development of lymphatic vessel valves (By similarity). Required to promote lymphatic

endothelial cell migration, in a process that involves down-regulation of integrin beta 1 activation and control of cell adhesion to the extracellular matrix (PubMed:<a href="http://www.uniprot.org/citations/35235341" target="\_blank">35235341</a>). Regulates the activity of mechanosensitive Piezo channel (PubMed:<a href="http://www.uniprot.org/citations/37590348" target="\_blank">37590348</a>).

#### Cellular Location

[Isoform 1]: Nucleus, nucleolus. Note=Also shows a granular distribution in the cytoplasm

#### Tissue Location


Expressed in lymphatic tissues. Detected in the spleen, thymus, peripheral blood leukocytes as well as prostate, uterus and small intestine. Expressed in lymphatic endothelial cells (PubMed:35235341).

### MDFIC Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### MDFIC Polyclonal Antibody - Images



#### Sample:


Liver (Mouse) Lysate at 40 ug

Primary: Anti-MDFIC (bs-6386R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 26 kD

Observed band size: 26 kD



**Sample:**

Liver (Rat) Lysate at 40 ug

Primary: Anti-MDFIC (bs-6386R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 26 kD

Observed band size: 26/27 kD