

CEE Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58438**Specification****CEE Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	Q7L5D6
Reactivity	Rat, Pig, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	36504

CEE Polyclonal Antibody - Additional Information**Gene ID** 51608**Other Names**

Golgi to ER traffic protein 4 homolog, Conserved edge-expressed protein, Transmembrane domain recognition complex 35 kDa subunit, TRC35, GET4 (http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=21690 target="_blank">HGNC:21690), C7orf20, CEE, TRC35

Dilution

WB~1:1000<br \>IHC-P~N/A<br \>IHC-F~N/A<br \>IF~1:50~200<br \>E~N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CEE Polyclonal Antibody - Protein Information**Name** GET4 ([HGNC:21690](#))**Synonyms** C7orf20, CEE, TRC35**Function**

As part of a cytosolic protein quality control complex, the BAG6/BAT3 complex, maintains misfolded and hydrophobic patches- containing proteins in a soluble state and participates in their proper delivery to the endoplasmic reticulum or alternatively can promote their sorting to the proteasome where they undergo degradation (PubMed:20676083, PubMed:<a

[21636303](http://www.uniprot.org/citations/21636303), PubMed: [21743475](http://www.uniprot.org/citations/21743475), PubMed: [28104892](http://www.uniprot.org/citations/28104892), PubMed: [32395830](http://www.uniprot.org/citations/32395830)). The BAG6/BAT3 complex is involved in the post-translational delivery of tail-anchored/type II transmembrane proteins to the endoplasmic reticulum membrane. Recruited to ribosomes, it interacts with the transmembrane region of newly synthesized tail-anchored proteins and together with SGTA and ASNA1 mediates their delivery to the endoplasmic reticulum (PubMed: [20676083](http://www.uniprot.org/citations/20676083), PubMed: [25535373](http://www.uniprot.org/citations/25535373), PubMed: [28104892](http://www.uniprot.org/citations/28104892)). Client proteins that cannot be properly delivered to the endoplasmic reticulum are ubiquitinated and sorted to the proteasome (PubMed: [28104892](http://www.uniprot.org/citations/28104892)). Similarly, the BAG6/BAT3 complex also functions as a sorting platform for proteins of the secretory pathway that are mislocalized to the cytosol either delivering them to the proteasome for degradation or to the endoplasmic reticulum (PubMed: [21743475](http://www.uniprot.org/citations/21743475)). The BAG6/BAT3 complex also plays a role in the endoplasmic reticulum-associated degradation (ERAD), a quality control mechanism that eliminates unwanted proteins of the endoplasmic reticulum through their retrotranslocation to the cytosol and their targeting to the proteasome. It maintains these retrotranslocated proteins in an unfolded yet soluble state condition in the cytosol to ensure their proper delivery to the proteasome (PubMed: [21636303](http://www.uniprot.org/citations/21636303)).

Cellular Location

Cytoplasm, cytosol

CEE Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CEE Polyclonal Antibody - Images