

SCN1B Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58489

Specification

SCN1B Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q07699
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	22 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human SCN1B
Epitope Specificity	21-120/268
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Membrane; Single-pass type I membrane protein.
SIMILARITY	Belongs to the sodium channel auxiliary subunit SCN1B (TC 8.A.17) family. Contains 1 Ig-like C2-type (immunoglobulin-like) domain.
SUBUNIT	The voltage-sensitive sodium channel consists of an ion conducting pore forming alpha-subunit regulated by one or more beta-1, beta-2 and beta-3. Beta-1 and beta-3 are non-covalently associated with alpha, while beta-2 is covalently linked by disulfide bonds. Beta-1 or beta-3 subunits associate with neurofascin. Associates with SCN10A (By similarity).
DISEASE	Defects in SCN1B are the cause of generalized epilepsy with febrile seizures plus type 1 (GEFS+1) [MIM:604233]. Generalized epilepsy with febrile seizures-plus refers to a rare autosomal dominant, familial condition with incomplete penetrance and large intrafamilial variability. Patients display febrile seizures persisting sometimes beyond the age of 6 years and/or a variety of afebrile seizure types. GEFS+ is a disease combining febrile seizures, generalized seizures often precipitated by fever at age 6 years or more, and partial

Important Note

seizures, with a variable degree of severity. Defects in SCN1B are the cause of Brugada syndrome type 5 (BRGDA5) [MIM:612838]. A tachyarrhythmia characterized by right bundle branch block and ST segment elevation on an electrocardiogram (ECG). It can cause the ventricles to beat so fast that the blood is prevented from circulating efficiently in the body. When this situation occurs (called ventricular fibrillation), the individual will faint and may die in a few minutes if the heart is not reset. This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Crucial in the assembly, expression, and functional modulation of the heterotrimeric complex of the sodium channel. The subunit beta-1 can modulate multiple alpha subunit isoforms from brain, skeletal muscle, and heart. Its association with neurofascin may target the sodium channels to the nodes of Ranvier of developing axons and retain these channels at the nodes in mature myelinated axons. Tissue specificity; Abundantly expressed in skeletal muscle, heart and brain.

SCN1B Polyclonal Antibody - Additional Information**Gene ID 6324****Other Names**

Sodium channel subunit beta-1, SCN1B

Target/Specificity

The overall expression of isoforms 1 and 2 is very similar. Isoform 1 is abundantly expressed in skeletal muscle, heart and brain. Isoform 2 is highly expressed in brain and skeletal muscle and present at a very low level in heart, placenta, lung, liver, kidney and pancreas. In brain, isoform 2 is most abundant in the cerebellum, followed by the cerebral cortex and occipital lobe, while isoform 1 levels are higher in the cortex compared to the cerebellum. Isoform 2 is expressed in many regions of the brain, including cerebellar Purkinje cells, cortex pyramidal neurons and many of the neuronal fibers throughout the brain (at protein level). Also detected in dorsal root ganglion, in fibers of the spinal nerve and in cortical neurons and their processes (at protein level).

Dilution

IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SCN1B Polyclonal Antibody - Protein Information**Name SCN1B ([HGNC:10586](#))**

Function

Regulatory subunit of multiple voltage-gated sodium (Nav) channels directly mediating the depolarization of excitable membranes. Navs, also called VGSCs (voltage-gated sodium channels) or VDSCs (voltage-dependent sodium channels), operate by switching between closed and open conformations depending on the voltage difference across the membrane. In the open conformation they allow Na⁺ ions to selectively pass through the pore, along their electrochemical gradient. The influx of Na⁺ ions provokes membrane depolarization, initiating the propagation of electrical signals throughout cells and tissues (PubMed:14622265, PubMed:15525788, PubMed:18464934, PubMed:19710327, PubMed:29992740, PubMed:36696443, PubMed:8125980, PubMed:8394762). The accessory beta subunits participate in localization and functional modulation of the Nav channels (PubMed:15525788, PubMed:19710327, PubMed:29992740). Modulates the activity of SCN1A/Nav1.1, SCN2A/Nav1.2, SCN3A/Nav1.3, SCN4A/Nav1.4, SCN5A/Nav1.5, SCN8A/Nav1.6, SCN9A/Nav1.7 and SCN10A/Nav1.8 (PubMed:14622265, PubMed:15525788, PubMed:18464934, PubMed:30765606, PubMed:36696443, PubMed:8125980, PubMed:8394762).

Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein. Perikaryon {ECO:0000250|UniProtKB:P97952} Cell projection {ECO:0000250|UniProtKB:P97952}. Cell projection, axon {ECO:0000250|UniProtKB:Q00954}. Note=Detected at nodes of Ranvier on the sciatic nerve. {ECO:0000250|UniProtKB:Q00954}

Tissue Location

The overall expression of isoform 1 and isoform 2 is very similar. Isoform 1 is abundantly expressed in skeletal muscle, heart and brain. Isoform 2 is highly expressed in brain and skeletal muscle and present at a very low level in heart, placenta, lung, liver, kidney and pancreas. In brain, isoform 2 is most abundant in the cerebellum, followed by the cerebral cortex and occipital lobe, while isoform 1 levels are higher in the cortex compared to the cerebellum. Isoform 2 is expressed in many regions of the brain, including cerebellar Purkinje cells, cortex pyramidal neurons and many of the neuronal fibers throughout the brain (at protein level). Also detected in dorsal root ganglion, in fibers of the spinal nerve and in cortical neurons and their processes (at protein level)

SCN1B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)

- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SCN1B Polyclonal Antibody - Images