

TULP2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58503

Specification

TULP2 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Bovine
Host
Clonality
Calculated MW
Physical State

O00295
Rabbit
Polyclonal
S9 KDa
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human TULP2

Epitope Specificity 301-400/520

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm. Secreted. Note=Does not have

a cleavable signal peptide and is secreted

by a non-conventional pathway

SIMILARITY Belongs to the TUB family.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

TULP2 is a member of a family of tubby-like genes (TULPs) that encode proteins of unknown function. Members of this family have been identified in plants, vertebrates, and invertebrates. The TULP proteins share a conserved C-terminal region of approximately 200 amino acid residues. [provided by RefSeq, Jul 2008]

TULP2 Polyclonal Antibody - Additional Information

Gene ID 7288

Other Names

Tubby-related protein 2, Cancer/testis antigen 65, CT65, Tubby-like protein 2, TULP2, TUBL2

Target/Specificity

Strongly expressed in testis. Also expressed in retina. Expressed in cancer cell lines.

Dilution

IHC-P~~N/A<br \><span class</pre>

="dilution_IHC-F">IHC-F~~N/A<br \><span class

="dilution IF">IF \sim 1:50 \sim 200<br\>E \sim N/A



Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

TULP2 Polyclonal Antibody - Protein Information

Name TULP2

Synonyms TUBL2

Cellular Location

Cytoplasm. Secreted. Note=Does not have a cleavable signal peptide and is secreted by a non-conventional pathway.

Tissue Location

Strongly expressed in testis. Also expressed in retina. Expressed in cancer cell lines.

TULP2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

TULP2 Polyclonal Antibody - Images