

Ubiquitously expressed transcript Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58504**Specification**

Ubiquitously expressed transcript Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	Q9UBK9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human UXT/Ubiquitously expressed transcript
Epitope Specificity	85-157/157
Isotype	IgG
Purity	
affinity purified by Protein A	
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SIMILARITY	Belongs to the UXT family.
SUBUNIT	Homohexamer. Interacts with LRPPRC. Interacts with AR N-terminus. In the nucleus interacts specifically with RELA and forms a dynamic complex with NF-kappa-B and is recruited to the NF-kappa-B enhanceosome upon stimulation. Interacts with MECOM and URI1.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

UXT is a novel protein, highly conserved in mouse. It interacts with the N terminus of the androgen receptor and plays a role in facilitating receptor-induced transcriptional activation. It is abundantly expressed in tumor tissues and is likely to be involved in tumorigenesis.

Ubiquitously expressed transcript Polyclonal Antibody - Additional Information**Gene ID** 8409**Other Names**

Protein UXT, Androgen receptor trapped clone 27 protein, ART-27, Ubiquitously expressed transcript protein, UXT

Target/Specificity

Ubiquitous. Expressed in prostate epithelial cells. Overexpressed in a number of tumor tissues (PubMed:16221885). Highest levels in the heart, skeletal muscle, pancreas, kidney, liver, adrenal gland, peripheral blood leukocytes, lymph node, prostate, and thyroid and the lowest levels in bladder and uterus.

Dilution

IHC-P ~ N/A
IHC-F ~ N/A
IF ~ 1:50 ~ 200
E ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glycerol

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Ubiquitously expressed transcript Polyclonal Antibody - Protein Information**Name** UXT**Function**

Involved in gene transcription regulation (PubMed:[21730289](http://www.uniprot.org/citations/21730289), PubMed:[28106301](http://www.uniprot.org/citations/28106301)). Acts in concert with the corepressor URI1 to regulate androgen receptor AR-mediated transcription (PubMed:[11854421](http://www.uniprot.org/citations/11854421), PubMed:[21730289](http://www.uniprot.org/citations/21730289)). Together with URI1, associates with chromatin to the NKX3-1 promoter region (PubMed:[21730289](http://www.uniprot.org/citations/21730289)). Negatively regulates the transcriptional activity of the estrogen receptor ESR1 by inducing its translocation into the cytoplasm (PubMed:[28106301](http://www.uniprot.org/citations/28106301)). May act as nuclear chaperone that facilitates the formation of the NF-kappa-B enhanceosome and thus positively regulates NF-kappa-B transcription activity (PubMed:[17620405](http://www.uniprot.org/citations/17620405), PubMed:[21307340](http://www.uniprot.org/citations/21307340)). Potential component of mitochondrial-associated LRPPRC, a multidomain organizer that potentially integrates mitochondria and the microtubular cytoskeleton with chromosome remodeling (PubMed:[17554592](http://www.uniprot.org/citations/17554592)). Increasing concentrations of UXT contributes to progressive aggregation of mitochondria and cell death potentially through its association with LRPPRC (PubMed:[17554592](http://www.uniprot.org/citations/17554592)). Suppresses cell transformation and it might mediate this function by interaction and inhibition of the biological activity of cell proliferation and survival stimulatory factors like MECOM (PubMed:[17635584](http://www.uniprot.org/citations/17635584)).

Cellular Location

[Isoform 1]: Cytoplasm Cytoplasm. Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole Note=Predominantly localizes to the nucleus (PubMed:16221885) Localizes to spindle pole during mitosis (PubMed:16221885)

Tissue Location

Ubiquitous (PubMed:10087202, PubMed:11827465, PubMed:11854421, PubMed:17635584). Expressed in prostate epithelial cells (PubMed:21730289). Expressed in mammary epithelial cells (PubMed:28106301). Highest levels in the heart, skeletal muscle, pancreas, kidney, liver, adrenal


gland, peripheral blood leukocytes, lymph node, prostate, and thyroid and the lowest levels in bladder and uterus (PubMed:11827465, PubMed:11854421, PubMed:17635584) Overexpressed in a number of tumor tissues (PubMed:11854421, PubMed:16221885, PubMed:28106301).

Ubiquitously expressed transcript Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Ubiquitously expressed transcript Polyclonal Antibody - Images




Paraformaldehyde-fixed, paraffin embedded (human liver carcinoma); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Protein UXT) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (Mouse stomach); Antigen retrieval by boiling in


sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse kidney); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Mouse intestine); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (Ubiquitously expressed transcript) Polyclonal Antibody, Unconjugated (bs-6749R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.