

BCA2/ZNF364 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58507

Specification

BCA2/ZNF364 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF, E

Primary Accession <u>O9Y4L5</u>

Reactivity
Host
Clonality
Rat, Pig, Bovine
Rabbit
Polyclonal

Clonality Polyclo
Calculated MW 34 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human ZNF364/BCA2/RNF115

Epitope Specificity 161-260/304

Isotype IgG
Purity

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm, cytosol (By similarity).

Note=The GTP-bound form of RAB7A recruits RNF115 from the cytosol onto late

endosomes/lysosomes (By similarity).

SIMILARITY Contains 1 RING-type zinc finger.

SUBUNIT Interacts with RAB7A.

Post-translational modifications RING-type zinc finger-dependent and

E2-dependent autoubiquitination.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

Background Descriptions

affinity purified by Protein A

The RING finger family of proteins possess ubiquitin ligase activity and play pivotal roles in protein degradation and receptor-mediated endocytosis. The autoubiquitination activity of ZNF364 indicates that it is a novel RING-type E3 ligase. Its effects on cell growth indicate that ZNF364 may be important for the ubiquitin modification of proteins crucial to breast carcinogenesis and growth.

BCA2/ZNF364 Polyclonal Antibody - Additional Information

Gene ID 27246

Other Names

E3 ubiquitin-protein ligase RNF115, 2.3.2.27, RING finger protein 115 {ECO:0000312|HGNC:HGNC:18154}, RING-type E3 ubiquitin transferase RNF115, Rabring 7, RNF115 (HGNC:18154)



Target/Specificity

Expressed at extremely low levels in normal breast, prostate, lung, colon. Higher levels of expression are detected in heart, skeletal muscle, testis as well as in breast and prostate cancer cells.

Dilution

WB~~1:1000<br \> <span class
="dilution_IHC-P">IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

BCA2/ZNF364 Polyclonal Antibody - Protein Information

Name RNF115 (HGNC:18154)

Function

E3 ubiquitin-protein ligase that catalyzes the 'Lys- 48'- and/or 'Lys-63'-linked polyubiquitination of various substrates and thereby plays a role in a number of signaling pathways including autophagy, innate immunity, cell proliferation and cell death (PubMed:20019814, PubMed:30689267). Plays a role in the endosomal trafficking and degradation of membrane receptors including EGFR, FLT3, MET and CXCR4 through their polyubiquitination. Participates together with BST2 in antiviral immunity by facilitating the internalization of HIV-1 virions into intracellular vesicles leading to their lysosomal degradation (PubMed: 20019814). Also possesses an antiviral activity independently of BST2 by promoting retroviral GAG proteins ubiquitination, redistribution to endo-lysosomal compartments and, ultimately, lysosomal degradation (PubMed:24852021). Catalyzes distinct types of ubiquitination on MAVS and STING1 at different phases of viral infection to promote innate antiviral response (PubMed:33139700). Mediates the 'Lys-48'-linked ubiquitination of MAVS leading to its proteasomal degradation and ubiquitinates STING1 via 'Lys-63'-linked polyubiquitination, critical for its oligomerization and the subsequent recruitment of TBK1 (PubMed: 33139700). Plays a positive role in the autophagosome-lysosome fusion by interacting with STX17 and enhancing its stability without affecting 'Lys-48'- or 'Lys-63'-linked polyubiquitination levels, which in turn promotes autophagosome maturation (PubMed:32980859). Negatively regulates TLR-induced expression of proinflammatory cytokines by catalyzing 'Lys-11'-linked ubiquitination of RAB1A and RAB13 to inhibit post-ER trafficking of TLRs to the Golgi by RAB1A and subsequently from the Golgi apparatus to the cell surface by RAB13 (PubMed:<a

Cellular Location

Cytoplasm. Nucleus Endoplasmic reticulum. Golgi apparatus. Note=The GTP-bound form of RAB7A recruits RNF115 from the cytosol onto late endosomes/lysosomes

href="http://www.uniprot.org/citations/35343654" target="_blank">35343654).

Tissue Location

Expressed at extremely low levels in normal breast, prostate, lung, colon. Higher levels of expression are detected in heart, skeletal muscle, testis as well as in breast and prostate cancer



cells.

BCA2/ZNF364 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BCA2/ZNF364 Polyclonal Antibody - Images