

**ILP2 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58538****Specification**

---

**ILP2 Polyclonal Antibody - Product Information**

Application	<b>IHC-P</b>
Primary Accession	<a href="#">Q96P09</a>
Reactivity	<b>Rat</b>
Host	<b>Rabbit</b>
Clonality	<b>Polyclonal</b>
Calculated MW	<b>27089</b>

**ILP2 Polyclonal Antibody - Additional Information****Other Names**

Baculoviral IAP repeat-containing protein 8, Inhibitor of apoptosis-like protein 2, IAP-like protein 2, ILP-2, Testis-specific inhibitor of apoptosis, BIRC8, ILP2

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ILP2 Polyclonal Antibody - Protein Information**

**Name** BIRC8

**Synonyms** ILP2

**Function**

Protects against apoptosis mediated by BAX.

**Cellular Location**

Cytoplasm.

**Tissue Location**

Testis specific in normal tissues.


**ILP2 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ILP2 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (rat brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (BIRC8) Polyclonal Antibody, Unconjugated (bs-6870R) at 1:400 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.