

**CCDC69 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58549****Specification**

---

**CCDC69 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF
Primary Accession	<a href="#">A6NI79</a>
Reactivity	Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	34796

**CCDC69 Polyclonal Antibody - Additional Information****Gene ID** 26112**Other Names**

Coiled-coil domain-containing protein 69, CCDC69 (<a href="http://www.genenames.org/cgi-bin/gene\_symbol\_report?hgnc\_id=24487" target="\_blank">HGNC:24487</a>)

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CCDC69 Polyclonal Antibody - Protein Information****Name** CCDC69 ([HGNC:24487](#))**Function**

May act as a scaffold to regulate the recruitment and assembly of spindle midzone components. Required for the localization of AURKB and PLK1 to the spindle midzone.

**Cellular Location**

Cytoplasm, cytoskeleton, spindle. Midbody Note=During early anaphase, localizes along overlapping interpolar microtubules between the separating chromosomes. During late anaphase, localizes to the center of spindle midzone. Concentrated at the midbody during telophase.

**Tissue Location**

Highly expressed in duodenum, esophagus, pancreas, prostate, salivary gland, thymus and urinary bladder

**CCDC69 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CCDC69 Polyclonal Antibody - Images**