

**ZNF521 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58562****Specification**

---

**ZNF521 Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | IHC-P, IHC-F, IF, E    |
| Primary Accession | <a href="#">Q96K83</a> |
| Reactivity        | Rat, Pig, Dog, Bovine  |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |
| Calculated MW     | 147866                 |

**ZNF521 Polyclonal Antibody - Additional Information****Gene ID** 25925**Other Names**

Zinc finger protein 521, Early hematopoietic zinc finger protein, LYST-interacting protein 3, ZNF521, EHZF, LIP3

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ZNF521 Polyclonal Antibody - Protein Information****Name** ZNF521**Synonyms** EHZF, LIP3**Function**

Transcription factor that can both act as an activator or a repressor depending on the context. Involved in BMP signaling and in the regulation of the immature compartment of the hematopoietic system. Associates with SMADs in response to BMP2 leading to activate transcription of BMP target genes. Acts as a transcriptional repressor via its interaction with EBF1, a transcription factor involved specification of B-cell lineage; this interaction preventing EBF1 to bind DNA and activate target genes.

**Cellular Location**

Nucleus.

#### **Tissue Location**


Predominantly expressed in hematopoietic cells. Present in organs and tissues that contain stem and progenitor cells, myeloid and/or lymphoid: placenta, spleen, lymph nodes, thymus, bone marrow and fetal liver. Within the hematopoietic system, it is abundant in CD34(+) cells but undetectable in mature peripheral blood leukocytes, and its levels rapidly decrease during the differentiation of CD34(+) cells in response to hemopoietins

#### **ZNF521 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **ZNF521 Polyclonal Antibody - Images**



Tissue/cell: rat spleen tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-EHZF/ZNF521 Polyclonal Antibody, Unconjugated(bs-6978R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining