

**CLCN3 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58563****Specification**

---

**CLCN3 Polyclonal Antibody - Product Information**

Application	IHC-P
Primary Accession	<a href="#">P51790</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	90966

**CLCN3 Polyclonal Antibody - Additional Information****Gene ID** 1182**Other Names**

H(+)/Cl(-) exchange transporter 3, Chloride channel protein 3, CLC-3, Chloride transporter CLC-3, CLCN3

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**CLCN3 Polyclonal Antibody - Protein Information****Name** CLCN3**Function**

[Isoform 1]: Strongly outwardly rectifying, electrogenic H(+)/Cl(-)exchanger which mediates the exchange of chloride ions against protons (By similarity). The CLC channel family contains both chloride channels and proton-coupled anion transporters that exchange chloride or another anion for protons (PubMed:<a href="http://www.uniprot.org/citations/29845874" target="\_blank">29845874</a>). The presence of conserved gating glutamate residues is typical for family members that function as antiporters (PubMed:<a href="http://www.uniprot.org/citations/29845874" target="\_blank">29845874</a>).

**Cellular Location**

[Isoform 1]: Early endosome membrane; Multi-pass membrane protein. Late endosome membrane; Multi-pass membrane protein. Lysosome membrane {ECO:0000250|UniProtKB:P51791}; Multi-pass membrane protein. Cell membrane {ECO:0000250|UniProtKB:P51792}; Multi-pass membrane protein. Note=Isoform 1 is localized mainly in late endosomes.


**Tissue Location**

Expressed primarily in tissues derived from neuroectoderm. Within the brain, its expression is particularly evident in the hippocampus, olfactory cortex, and olfactory bulb. Highly expressed in aortic and coronary vascular smooth muscle cells, and aortic endothelial cells. Also expressed in tracheal and alveolar epithelial cells, and intima and media of the pulmonary vessels. Expressed in bronchus and colon (at protein level)

**CLCN3 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**CLCN3 Polyclonal Antibody - Images**

Tissue/cell: rat brain tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer ( 0.01M, pH 6.0 ), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-CLCN3/CLC-3 Polyclonal Antibody, Unconjugated(bs-6981R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CLCN3) Polyclonal Antibody, Unconjugated (bs-6981R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.