

**STEAP4 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58574****Specification****STEAP4 Polyclonal Antibody - Product Information**

Application	IHC-P, IHC-F, IF, E
Primary Accession	<a href="#">Q687X5</a>
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	51981

**STEAP4 Polyclonal Antibody - Additional Information****Gene ID** 79689**Other Names**

Metalloreductase STEAP4, 1.16.1.9, Six-transmembrane epithelial antigen of prostate 4, SixTransMembrane protein of prostate 2, Tumor necrosis factor, alpha-induced protein 9, STEAP4, STAMP2 {ECO:0000303|PubMed:15897894}, TNFAIP9

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
E ~ ~ N/A

**Format**

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**STEAP4 Polyclonal Antibody - Protein Information****Name** STEAP4**Synonyms** STAMP2 {ECO:0000303|PubMed:15897894}, TN**Function**

Integral membrane protein that functions as a NADPH-dependent ferric-chelate reductase, using NADPH from one side of the membrane to reduce a Fe(3+) chelate that is bound on the other side of the membrane. Mediates sequential transmembrane electron transfer from NADPH to FAD and onto heme, and finally to the Fe(3+) chelate (PubMed:[30337524](http://www.uniprot.org/citations/30337524)). Can also reduce Cu(2+) to Cu(1+) (By similarity). Plays a role in systemic metabolic homeostasis, integrating inflammatory and metabolic responses (By similarity). Associated with obesity and

insulin-resistance (PubMed:<a href="http://www.uniprot.org/citations/18381574" target="\_blank">18381574</a>, PubMed:<a href="http://www.uniprot.org/citations/18430367" target="\_blank">18430367</a>). Involved in inflammatory arthritis, through the regulation of inflammatory cytokines (PubMed:<a href="http://www.uniprot.org/citations/19660107" target="\_blank">19660107</a>). Inhibits anchorage- independent cell proliferation (PubMed:<a href="http://www.uniprot.org/citations/19787193" target="\_blank">19787193</a>).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein

#### **Tissue Location**

Ubiquitous. Highly expressed in adipose tissue. Expressed in placenta, lung, heart and prostate. Detected at lower levels in liver, skeletal muscle, pancreas, testis and small intestine Highly expressed in joints of patients with rheumatoid arthritis and localized with CD68 cells, a marker for macrophages

### **STEAP4 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **STEAP4 Polyclonal Antibody - Images**