

CDC2L6 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58585**Specification**

CDC2L6 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF, E
Primary Accession	O9BWU1
Reactivity	Rat, Chimpanzee, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	56802

CDC2L6 Polyclonal Antibody - Additional Information**Gene ID** 23097**Other Names**

Cyclin-dependent kinase 19, 2.7.11.22, CDC2-related protein kinase 6, Cell division cycle 2-like protein kinase 6, Cell division protein kinase 19, Cyclin-dependent kinase 11, Death-preventing kinase, CDK19, CDC2L6, CDK11, KIAA1028

Dilution

IHC-P ~ ~ N/A
IHC-F ~ ~ N/A
IF ~ ~ 1:50 ~ 200
E ~ ~ N/A

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

CDC2L6 Polyclonal Antibody - Protein Information**Name** CDK19**Synonyms** CDC2L6, CDK11, KIAA1028**Cellular Location**


Cytoplasm. Cytoplasm, perinuclear region. Nucleus

CDC2L6 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CDC2L6 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (CDC2L6) Polyclonal Antibody, Unconjugated (bs-7070R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.