

**ZNF229 Polyclonal Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP58628**

**Specification**

**ZNF229 Polyclonal Antibody - Product Information**

|                                |   |
|--------------------------------|---|
| Application                    | IHC-P, IHC-F, IF, ICC, E  |
| Primary Accession              | <a href="#">O9UJW7</a>  |
| Host                           | Rabbit  |
| Clonality                      | Polyclonal  |
| Calculated MW                  | 94 KDa  |
| Physical State                 | Liquid  |
| Immunogen                      | KLH conjugated synthetic peptide derived from human ZNF229  |
| Epitope Specificity            | 201-300/825   |
| Isotype                        | IgG   |
| <b>Purity</b>                  |   |
| affinity purified by Protein A |   |
| Buffer                         | 0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.   |
| SUBCELLULAR LOCATION           | Nucleus.  |
| SIMILARITY                     | Belongs to the krueppel C2H2-type zinc-finger protein family. Contains 18 C2H2-type zinc fingers. Contains 1 KRAB domain. |
| Important Note                 | This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications. |

**ZNF229 Polyclonal Antibody - Additional Information**

**Gene ID** 7772

**Other Names**

Zinc finger protein 229, ZNF229

**Dilution**

IHC-P ~ ~ N/A  
IHC-F ~ ~ N/A  
IF ~ ~ 1:50 ~ 200  
ICC ~ ~ N/A  
E ~ ~ N/A

**Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

**ZNF229 Polyclonal Antibody - Protein Information**

**Name** ZNF229 ([HGNC:13022](#))

**Function**

May be involved in transcriptional regulation.

**Cellular Location**


Nucleus.

**ZNF229 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**ZNF229 Polyclonal Antibody - Images**



Paraformaldehyde-fixed, paraffin embedded (human liver); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ZNF229) Polyclonal Antibody, Unconjugated (bs-7189R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.