

RhoGAP5 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58657

Specification

RhoGAP5 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, ICC, E

Primary Accession Q13017

Reactivity Rat, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 172 KDa
Physical State Liquid

Immunogen KLH conjugated synthetic peptide derived

from human RhoGAP5

Epitope Specificity 251-350/1502

Isotype IgG

Purityaffinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasmic.

SIMILARITY Contains 4 FF domains. Contains 1

Rho-GAP domain.

Important Note This product as supplied is intended for

research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Rho GTPase activating protein 5 negatively regulates RHO GTPases, a family which may mediate cytoskeleton changes by stimulating the hydrolysis of bound GTP. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

RhoGAP5 Polyclonal Antibody - Additional Information

Gene ID 394

Other Names

Rho GTPase-activating protein 5, Rho-type GTPase-activating protein 5, p190-B, ARHGAP5, RHOGAP5

Target/Specificity

Expressed in kidney, brain, liver and lung.

Dilution

IHC-P~~N/A<br \><span class</pre>

="dilution_IHC-F">IHC-F~~N/A<br \><span class

="dilution_IF">IF \sim 1:50 \sim 200<br \>ICC \sim N/A<br \>E \sim N/A



Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RhoGAP5 Polyclonal Antibody - Protein Information

Name ARHGAP5

Synonyms RHOGAP5

Function

GTPase-activating protein for Rho family members (PubMed:8537347).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Note=Also membrane-associated in a fibrillar pattern that colocalizes with the alpha5-beta1 integrin receptor (ITGA5/ITGB1) for fibronectin.

Tissue Location

Detected in skin fibroblasts (at protein level) (PubMed:8537347).

RhoGAP5 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

RhoGAP5 Polyclonal Antibody - Images