

PSRC1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58693

Specification

PSRC1 Polyclonal Antibody - Product Information

Application WB, IHC-P, IHC-F, IF

Primary Accession <u>Q6PGN9</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 38796

PSRC1 Polyclonal Antibody - Additional Information

Gene ID 84722

Other Names

Proline/serine-rich coiled-coil protein 1, PSRC1, DDA3

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

PSRC1 Polyclonal Antibody - Protein Information

Name PSRC1

Synonyms DDA3

Function

Required for normal progression through mitosis. Required for normal congress of chromosomes at the metaphase plate, and for normal rate of chromosomal segregation during anaphase. Plays a role in the regulation of mitotic spindle dynamics. Increases the rate of turnover of microtubules on metaphase spindles, and contributes to the generation of normal tension across sister kinetochores. Recruits KIF2A and ANKRD53 to the mitotic spindle and spindle poles. May participate in p53/TP53-regulated growth suppression.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, spindle pole. Note=Detected at the mitotic spindle and spindle poles. Diffusely distributed throughout the cell during interphase

Tissue Location

Widely expressed in adult and fetal tissues, with highest expression in the adult brain and fetal



thymus. Not detected in adult skeletal muscle.

PSRC1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

PSRC1 Polyclonal Antibody - Images