

C4orf32 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5870c**Specification**

C4orf32 Antibody (Center) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q8N8J7
Other Accession	NP_689613.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	14653
Antigen Region	30-56

C4orf32 Antibody (Center) - Additional Information**Gene ID** 132720**Other Names**

Uncharacterized protein C4orf32, C4orf32

Target/Specificity

This C4orf32 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 30-56 amino acids from the Central region of human C4orf32.

Dilution

WB~~1:1000

IHC-P~~1:50~100

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

C4orf32 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

C4orf32 Antibody (Center) - Protein Information**Name** FAM241A ([HGNC:26813](#))**Synonyms** C4orf32


Cellular Location

Membrane; Single-pass membrane protein


C4orf32 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

C4orf32 Antibody (Center) - Images

C4orf32 Antibody (Center) (Cat. #AP5870c) western blot analysis in Ramos cell line lysates (35ug/lane). This demonstrates the C4orf32 antibody detected the C4orf32 protein (arrow).



C4orf32 antibody (Center) (Cat. #AP5870c) immunohistochemistry analysis in formalin fixed and paraffin embedded human lymph tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the C4orf32 antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.