

BCCIP Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58712

Specification

BCCIP Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E

Primary Accession <u>O9P287</u>

Reactivity Rat, Pig, Dog, Bovine

Host Rabbit
Clonality Polyclonal
Calculated MW 35979

BCCIP Polyclonal Antibody - Additional Information

Gene ID 56647

Other Names

BRCA2 and CDKN1A-interacting protein, P21- and CDK-associated protein 1, Protein TOK-1, BCCIP, TOK1

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

BCCIP Polyclonal Antibody - Protein Information

Name BCCIP


Synonyms TOK1

Function

During interphase, required for microtubule organizing and anchoring activities. During mitosis, required for the organization and stabilization of the spindle pole (PubMed:28394342). Isoform 2/alpha is particularly important for the regulation of microtubule anchoring, microtubule stability, spindle architecture and spindle orientation, compared to isoform 1/beta (PubMed:28394342). May promote cell cycle arrest by enhancing the inhibition of CDK2 activity by CDKN1A. May be required for repair of DNA damage by homologous recombination in conjunction with BRCA2. May not be involved in non-homologous end joining (NHEJ).

Cellular Location







Nucleus. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriole. Cytoplasm, cytoskeleton, spindle pole. Note=Colocalizes with BRCA2 in discrete nuclear foci (PubMed:15713648). In interphase, preferential localizes to the mother centriole (PubMed:28394342). Recruited to the spindle pole matrix and centrosome by microtubules and dynein/dynactin activity (PubMed:28394342). [Isoform 2]: Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle pole. Note=Isoform 2/alpha tends to be more abundant at, and more strongly associated with, centrosomes than isoform 1/beta.

Tissue Location

Expressed at high levels in testis and skeletal muscle and at lower levels in brain, heart, kidney, liver, lung, ovary, pancreas, placenta, and spleen.

BCCIP Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

BCCIP Polyclonal Antibody - Images