

RBM38 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58715**Specification**

RBM38 Polyclonal Antibody - Product Information

Application	IHC-P, IHC-F, IF
Primary Accession	O9H0Z9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	25498

RBM38 Polyclonal Antibody - Additional Information**Gene ID** 55544**Other Names**

RNA-binding protein 38, CLL-associated antigen KW-5, HSRNASEB, RNA-binding motif protein 38, RNA-binding region-containing protein 1, ssDNA-binding protein SEB4, RBM38, RNPC1, SEB4

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RBM38 Polyclonal Antibody - Protein Information**Name** RBM38**Synonyms** RNPC1, SEB4**Function**

RNA-binding protein that specifically bind the 3'-UTR of CDKN1A transcripts, leading to maintain the stability of CDKN1A transcripts, thereby acting as a mediator of the p53/TP53 family to regulate CDKN1A. CDKN1A is a cyclin-dependent kinase inhibitor transcriptionally regulated by the p53/TP53 family to induce cell cycle arrest. Isoform 1, but not isoform 2, has the ability to induce cell cycle arrest in G1 and maintain the stability of CDKN1A transcripts induced by p53/TP53. Also acts as a mRNA splicing factor. Specifically regulates the expression of FGFR2-IIIb, an epithelial cell-specific isoform of FGFR2. Plays a role in myogenic differentiation.

Cellular Location

Cytoplasm, cytosol. Nucleus

RBM38 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBM38 Polyclonal Antibody - Images