

### **TRIP1 Polyclonal Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58735

### **Specification**

# **TRIP1 Polyclonal Antibody - Product Information**

Application WB, IHC-P, IHC-F, IF, E

Primary Accession
Reactivity
Rat
Host
Clonality
Calculated MW
Physical State
Reactivity
Rat
Rabbit
Polyclonal
Liquid

Immunogen KLH conjugated synthetic peptide derived

from human TRIP1/PSMC5

Epitope Specificity 122-320/406

Isotype IgG
Purity

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cytoplasm (Potential). Nucleus (Potential).

SIMILARITY
SUBUNIT

Belongs to the AAA ATPase family.
Interacts, in vitro, with the thyroid
hormone receptor (in a thyroid hormone
T3-dependent manner) and with retinoid X
receptor (RXR) (By similarity). Interacts
with NDC80. Interacts with PAAF1.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

### **Background Descriptions**

Human SUG1 (also known as p45, Rpt6, Cim3 and PSMC5) is a component of the 19S regulatory subunit of the 26S proteosome complex. It is one of the AAA ATPas found in the 19S regulatory complex that is thought to participate in the unfolding of ubiquitinated proteins in an ATP dependent manner. It has also been shown to directly interact with the nucleotide excision repair protein XPB.

### **TRIP1** Polyclonal Antibody - Additional Information

# **Gene ID 5705**

# **Other Names**

26S proteasome regulatory subunit 8, 26S proteasome AAA-ATPase subunit RPT6, Proteasome 26S subunit ATPase 5, Proteasome subunit p45, Thyroid hormone receptor-interacting protein 1, TRIP1, p45/SUG, PSMC5, SUG1

#### **Dilution**



<span class="dilution_WB">WB~~1:1000</span> <span class<="" th=""></span>
="dilution_IHC-P">IHC-P~~N/A <span class<="" td=""></span>
="dilution_IHC-F">IHC-F~~N/A <span class<="" td=""></span>
="dilution $IF">IF~\sim 1:50\sim 200or \>E\sim\sim N/A$

### **Storage**

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **TRIP1 Polyclonal Antibody - Protein Information**

Name PSMC5

Synonyms SUG1

### **Function**

Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair. PSMC5 belongs to the heterohexameric ring of AAA (ATPases associated with diverse cellular activities) proteins that unfolds ubiquitinated target proteins that are concurrently translocated into a proteolytic chamber and degraded into peptides.

**Cellular Location** 

Cytoplasm. Nucleus.


# **TRIP1 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **TRIP1 Polyclonal Antibody - Images**






Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIP1) Polyclonal Antibody, Unconjugated (bs-7731R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.




Paraformaldehyde-fixed, paraffin embedded (rat lung); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (TRIP1) Polyclonal Antibody, Unconjugated (bs-7731R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.



Sample:

Kidney (Mouse) Lysate at 40 ug

Primary: Anti-TRIP1 (bs-7731R) at 1/1000 dilution





Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 45 kD Observed band size: 45 kD