

SPAG8 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP58786

Specification

SPAG8 Polyclonal Antibody - Product Information

Application IHC-P, IHC-F, IF, E
Primary Accession Q99932

Reactivity
Rost
Clonality
Calculated MW
Clossion

Rat, Pig
Rabbit
Polyclonal
51139

SPAG8 Polyclonal Antibody - Additional Information

Gene ID 26206

Other Names

Sperm-associated antigen 8, HSD-1, Sperm membrane protein 1, SMP-1, Sperm membrane protein BS-84, SPAG8

Dilution

IHC-P~~N/A<br \> IHC-F~~N/A<br \> IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SPAG8 Polyclonal Antibody - Protein Information

Name SPAG8 (HGNC:14105)

Function

Microtubule inner protein (MIP) part of the dynein-decorated doublet microtubules (DMTs) in cilia axoneme, which is required for motile cilia beating (PubMed:36191189). Plays a role in spermatogenesis by enhancing the binding of CREM isoform tau to its coactivator FHL5 and increasing the FHL5-regulated transcriptional activation of CREM isoform tau (By similarity). Involved in the acrosome reaction and in binding of sperm to the zona pellucida (By similarity). Plays a role in regulation of the cell cycle by controlling progression through the G2/M phase, possibly by delaying the activation of CDK1 which is required for entry into mitosis (PubMed:19548270). May play a role in fertility and microtubule formation through interaction with RANBP9 (PubMed:10500252/a>).

Cellular Location



Tel: 858.875.1900 Fax: 858.875.1999

Cytoplasm {ECO:0000250|UniProtKB:Q3V0Q6}. Nucleus {ECO:0000250|UniProtKB:Q3V0Q6}. Cytoplasmic vesicle, secretory vesicle, acrosome. Cytoplasm, cytoskeleton, microtubule organizing center. Cytoplasm, cytoskeleton, spindle. Cytoplasm, cytoskeleton, cilium axoneme. Cytoplasm, cytoskeleton, flagellum axoneme {ECO:0000250|UniProtKB:Q3V0Q6}. Note=In mature sperm cells, detected in the acrosomal region of the head and in the middle piece of the tail (By similarity). Localized to the nucleus and cytoplasm of spermatocytes and round spermatids while, in elongating spermatids, expressed in the cytoplasm but not in the nucleus (By similarity). During the cell cycle, localized on the microtubule-organizing center (MTOC) during prophase. In metaphase, extends along spindle microtubules. In anaphase, detected on the astral microtubules and mid-zone. In telophase, remains at the mid-zone. After cytokinesis, returns to the MTOC (PubMed:19548270). Microtubule inner protein (MIP) part of the dynein-decorated doublet microtubules (DMTs) in cilia axoneme (By similarity). {ECO:0000250|UniProtKB:E1BNS6, ECO:0000250|UniProtKB:Q3V0Q6, ECO:0000269|PubMed:19548270}

Tissue Location

Expressed in testis (germ cells), but not in liver, kidney, prostate and small intestine. Expressed in airway epithelial cells (PubMed:36191189).

SPAG8 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

SPAG8 Polyclonal Antibody - Images