

HOXC9 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP58821**Specification****HOXC9 Polyclonal Antibody - Product Information**

Application	WB, IHC-P, IHC-F, IF, E
Primary Accession	P31274
Reactivity	Rat, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	29 KDa
Physical State	Liquid
Immunogen	KLH conjugated synthetic peptide derived from human HOXC9
Epitope Specificity	151-220/260
Isotype	IgG
Purity	affinity purified by Protein A
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Nucleus.
SIMILARITY	Belongs to the Abd-B homeobox family. Contains 1 homeobox DNA-binding domain.
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

HOXC9 plays an important role in morphogenesis in all multicellular organisms. Mammals possess four similar homeobox gene clusters, HOXA, HOXB, HOXC and HOXD, which are located on different chromosomes and consist of 9 to 11 genes arranged in tandem.

HOXC9 Polyclonal Antibody - Additional Information**Gene ID 3225****Other Names**

Homeobox protein Hox-C9, Homeobox protein Hox-3B, HOXC9, HOX3B

Dilution

WB~~1:1000
IHC-P~~N/A
IHC-F~~N/A
IF~~1:50~200
E~~N/A

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH

7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

HOXC9 Polyclonal Antibody - Protein Information

Name HOXC9

Synonyms HOX3B

Function

Sequence-specific transcription factor which is part of a developmental regulatory system that provides cells with specific positional identities on the anterior-posterior axis.

Cellular Location


Nucleus.

HOXC9 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


HOXC9 Polyclonal Antibody - Images



Tissue/cell: human kidney tissue; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-HOXC9 Polyclonal Antibody, Unconjugated(bs-7982R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining

**Sample:**

Lane 1: Muscle (Mouse) Tissue Lysate at 40 ug
Lane 2: Muscle (Rat) Tissue Lysate at 40 ug
Lane 3: Ear (Rat) Tissue Lysate at 40 ug
Primary: Anti-HOXC9 (bs-7982R) at 1/200 dilution
Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution
Predicted band size: 29 kD
Observed band size: 40 kD